

EXISTENCE OF (N, λ) -PERIODIC SOLUTIONS FOR ABSTRACT FRACTIONAL DIFFERENCE EQUATIONS

EDGARDO ALVAREZ, STIVEN DÍAZ, AND CARLOS LIZAMA

ABSTRACT. We establish sufficient conditions for the existence and uniqueness of (N, λ) -periodic solutions for the following abstract model

$$\Delta^\alpha u(n) = Au(n+1) + f(n, u(n)), \quad n \in \mathbb{Z},$$

where $0 < \alpha \leq 1$, A is a closed linear operator defined in a Banach space X , Δ^α denotes the fractional difference operator in the Weyl-like sense and f satisfies appropriate conditions.

1. INTRODUCTION

In this article, we investigate the existence of a class of solutions for the abstract fractional difference equation

$$(1.1) \quad \Delta^\alpha u(n) = Au(n+1) + f(n, u(n)), \quad n \in \mathbb{Z},$$

called (N, λ) -periodic solutions. In (1.1), A is a possibly unbounded operator defined on a Banach space X and $f : \mathbb{Z} \times X \rightarrow X$ is given. This class of (N, λ) -periodic functions was introduced in the reference [6] as the discrete counterpart of the notion of (ω, c) -periodic functions [10], a notion that has been studied by various authors, see e.g. [8, 9, 15, 16, 17, 18, 20, 26] and [30]. It is worth noting that class of (N, λ) -periodic functions contains the classes of discrete periodic ($\lambda = 1$), discrete anti-periodic ($\lambda = -1$), discrete Bloch-periodic ($\lambda = e^{ikN}$, $k \in \mathbb{Z}$ fixed) and unbounded functions.

Existence and uniqueness of (N, λ) -periodic solutions for *scalar models*, such as Volterra difference equations with infinite delay, were recently investigated in [6]. Anticipating a growing theoretical and practical interest in this class of solutions, we study in this article the existence and uniqueness of solutions for the abstract Cauchy problem (1.1). This problem extends the class of models under study so far, to the broader context of partial difference-differential equations, that is *vector-valued models*, since A could be an unbounded operator defined in a Banach space X , e.g. the Laplacian operator on $L^2(\mathbb{R}^d)$. In fact, as we will reveal in this investigation, when A is the generator of a C_0 -semigroup that is strictly contractive, plus other conditions on the non-linear term, then the existence of (N, λ) -periodic solutions for (1.1) can be guaranteed.

2010 *Mathematics Subject Classification.* 39A14; 39A12; 65Q10; 47D06; 47D60.

Key words and phrases. (N, λ) -periodic solutions; Banach space; fractional difference operator; subordination.

Discrete fractional calculus has received considerable interest in recent years due to its interaction with applied mathematics and computation. We refer to the articles [13] for applications to interval-valued systems, [3] for applications to chaotic systems with short memory and image encryption, [14] for applications to variable-order fractional discrete-time recurrent neural networks, [31] for applications to image enhancement and [32] for applications to Lyapunov functions for Riemann-Liouville-like fractional difference equations.

The existence of solutions for the abstract model (1.1) began to be studied in the articles [21] and [12] in its linearized form. Subsequently, maximal regularity in Lebesgue spaces of sequences was studied in [23]. In case A is bounded, weighted bounded solutions were studied in [24]. In [1] the existence of almost automorphic mild solutions was studied.

However, the existence and uniqueness of (N, λ) -periodic solutions is an open topic that deserves to be investigated. The objective of this work is to solve this problem.

As methods, we use the technique of *resolvent sequences of operators*, a tool that was introduced in 2017 by Lizama [21] and has been used in several articles since then. Using this method, an explicit representation of the solution for (1.1) can be obtained, which allows the use of several fixed point theorems.

Recently it has been shown that resolvent sequences of operators can be related to each other by means of a subordination principle [7], at least when $0 < \alpha \leq 1$. In this work, we will refine the results in [7] observing that the only necessary condition to obtain the existence of resolvent sequences of operators is: $1 \in \rho(A)$, the resolvent set of A . This crucial observation is stated in the Theorem 3.1 below, and follows from the subordination principle which we will generalize here by using an extension of the discrete version of the Lévy α -stable distribution.

So our main result regarding the solubility of (1.1) can be proved and it says the following: Suppose that $1 \in \rho(A)$ and

$$(1.2) \quad r_A := \|(I - A)^{-1}\| < 1.$$

Assume that there exists $(N, \lambda) \in \mathbb{N} \times (\mathbb{C} \setminus \mathbb{D})$ and a constant $L > 0$ such that $f(n + N, \lambda x) = \lambda f(n, x)$ for all $(n, x) \in \mathbb{Z} \times X$ and

$$\|f(n, x) - f(n, y)\| \leq L\|x - y\|,$$

for all $x, y \in X$ and all $n \in \mathbb{Z}$. If

$$(1.3) \quad L < \left(1 - \frac{1}{|\lambda|^{1/N}}\right)^\alpha + \left(\frac{1}{r_A} - 1\right),$$

then equation (1.1) has a unique (N, λ) -periodic solution in a mild sense.

This article is organized as follows: Section 2 is devoted to the preliminaries about the notion of fractional order difference operator Δ^α that we will use. We will also remember the notion of Mittag-Leffler sequence and of (μ, ν) -resolvent sequence of operators. We end this section by remembering the definition of (N, λ) -periodic

sequence. Section 3 introduces the notion of a scaled Wright sequence and lists some of its main properties. Also in this section we establish the important Theorem 3.1 showing that the condition $1 \in \rho(A)$ is sufficient for the existence of (μ, ν) -resolvent sequence of operators. Then we prove the Theorem 3.4 which says that under the condition (1.2) the summability of (ν, ν) -resolvent sequences can be ensured. Using this relevant fact, we solve in Section 4 the problem of existence and uniqueness of (N, λ) -periodic solutions for the equation (1.1). See Theorem 4.5. Finally, an example is given where A is the one-dimensional Laplacian in $X = L^2(0, 1)$.

2. PRELIMINARIES.

Let X be a complex Banach space with norm $\|\cdot\|$ and $\mathcal{B}(X)$ denotes the Banach space of all bounded operators defined on X . For a real number a , we denote $\mathbb{N}_a := \{a, a+1, a+2, \dots\}$ and when $a = 1$ we write \mathbb{N} . We recall that the finite discrete convolution $*$ of two sequences $f, g : \mathbb{N}_0 \rightarrow X$ is defined by

$$(f * g)(n) := \sum_{j=0}^n f(n-j)g(j), \quad n \in \mathbb{N}_0.$$

We denote by $s(\mathbb{Z}, X)$ the vector space consisting of all vector-valued sequences $f : \mathbb{Z} \rightarrow X$. For $f \in s(\mathbb{Z}, X)$, we recall that the forward difference operator $\Delta : s(\mathbb{Z}, X) \rightarrow s(\mathbb{Z}, X)$ is defined by

$$\Delta f(n) := f(n+1) - f(n), \quad n \in \mathbb{Z}.$$

On the other hand, for an arbitrary $\alpha \in \mathbb{C} \setminus \{0, -1, -2, \dots\}$ the Cesàro sequence $\{k^\alpha(n)\}_{n \in \mathbb{N}_0}$, introduced in [21] (see also [33]), is defined by

$$(2.1) \quad k^\alpha(n) := \frac{\Gamma(\alpha + n)}{\Gamma(\alpha)n!}, \quad n \in \mathbb{N}_0.$$

In case $\alpha = 0$ we define $k^0(n) := \delta_0(n)$ the Kronecker delta.

The following equality and estimate holds: for $\alpha > 0$

$$k^\alpha(n) = \frac{1}{n^{1-\alpha}\Gamma(\alpha)} \left(1 + O\left(\frac{1}{n}\right) \right)$$

and for $0 < \alpha < 1$,

$$(2.2) \quad \frac{1}{\Gamma(\alpha)(n+1)^{1-\alpha}} < k^\alpha(n) < \frac{1}{\Gamma(\alpha)n^{1-\alpha}}, \quad n \in \mathbb{N}.$$

Further, given $\alpha, \beta > 0$, the sequence k^α satisfies the semigroup property in \mathbb{N}_0 , that is,

$$(2.3) \quad (k^\alpha * k^\beta)(n) = k^{\alpha+\beta}(n), \quad \forall n \in \mathbb{N}_0,$$

see [2, Section 2]. Given $\alpha > 0$, we define the set

$$\ell_\alpha^1(\mathbb{Z}, X) := \left\{ f \in s(\mathbb{Z}, X) : \sum_{n=-\infty}^{\infty} \|n^{\alpha-1}f(n)\| < \infty \right\}.$$

It is clear that $\ell_\alpha^1(\mathbb{Z}, X)$ is a Banach space under the norm $\|f\|_{\ell_\alpha^1} := \sum_{n=-\infty}^{\infty} \|n^{\alpha-1}f(n)\|$.

If $\alpha = 1$ then we simply write $\ell^1(\mathbb{Z}, X)$. Now, suppose that $0 < \alpha \leq 1$. Observe that, if $f \in \ell^1(\mathbb{Z}, X)$ then

$$\|f\|_{\ell_\alpha^1} := \sum_{n=-\infty}^{\infty} \|n^{\alpha-1}f(n)\| < \sum_{n=-\infty}^{\infty} \|f(n)\| < \infty.$$

Hence, $\ell^1(\mathbb{Z}, X) \subset \ell_\alpha^1(\mathbb{Z}, X)$ for $0 < \alpha \leq 1$.

The theory and applications of operators defined by means of the Cesàro sequences defined on \mathbb{N}_0 have been worked in different investigations (see for example [2, 7, 11, 21, 22, 27]). In this paper we will work with the following fractional sum operator defined on \mathbb{Z} in the reference [1].

Definition 2.1 ([1]). *Given $0 < \alpha < 1$ the α -th fractional sum operator $\Delta^{-\alpha} : \ell^1(\mathbb{Z}, X) \rightarrow s(\mathbb{Z}, X)$ is defined by means of the formula*

$$(2.4) \quad \Delta^{-\alpha}f(n) := \sum_{j=-\infty}^n k^\alpha(n-j)f(j), \quad f \in \ell^1(\mathbb{Z}, X).$$

Remark 2.2. *Note that, for $\alpha = 0$, $\Delta^{-\alpha}f(n) = f(n)$.*

The next definition about fractional differences operators in the sense of Riemann-Liouville and Caputo was introduced by Abadias and Lizama in [1].

Definition 2.3. *Let $0 < \alpha < 1$ and $f \in \ell^1(\mathbb{Z}, X)$. The Caputo fractional difference operator of order α is defined by*

$${}_c\Delta^\alpha f(n) := \Delta^{-(1-\alpha)}\Delta f(n),$$

and the Riemann-Liouville fractional difference operator of order α is defined by

$$(2.5) \quad {}_R\Delta^\alpha f(n) := \Delta\Delta^{-(1-\alpha)}f(n).$$

Given $f \in \ell_\alpha^1(\mathbb{Z}, X)$, it was proved in [1] that

$${}_R\Delta^\alpha f(n) = {}_c\Delta^\alpha f(n), \quad n \in \mathbb{Z}.$$

Therefore, from now on, we will simply denote by Δ^α either ${}_R\Delta^\alpha$ or ${}_c\Delta^\alpha$.

Now, we recall the notion of *Mittag-Leffler sequence* defined and studied in the references [7, 21, 25, 27].

Let $\alpha, \beta > 0$ and $\sigma \in \mathbb{C}$ be such that $|\sigma| < 1$. We define

$$(2.6) \quad \mathcal{E}_{\alpha,\beta}(\sigma, n) := \sum_{j=0}^{\infty} \sigma^j k^{\alpha j + \beta}(n), \quad n \in \mathbb{N}_0.$$

Note that the series on the right hand side of (2.6) converges by (2.2). Furthermore, the Z -transform of the Mittag-Leffler sequence exists for $|z| > 1$ (see [7]), and is given by

$$(2.7) \quad \sum_{j=0}^{\infty} \mathcal{E}_{\alpha, \beta}(\sigma, j) z^{-j} = \left(\frac{z}{z-1} \right)^{\beta} \left(1 - \sigma \left(\frac{z}{z-1} \right)^{\alpha} \right)^{-1}.$$

Motivated by [7, (the proof of) Proposition 4.6], we get the following result regarding the asymptotic behavior of the Mittag-Leffler sequence.

Theorem 2.4. *Let $0 < \alpha < 2$, $\beta, C > 0$ and $\sigma \in \mathbb{C}$ be such that $|\sigma| < 1$. The Mittag-Leffler sequence (2.6) satisfies the following inequality*

$$|\mathcal{E}_{\alpha, \beta}(-\sigma, n)| \leq \frac{C}{1 + \sigma n^{\alpha}}, \quad n \in \mathbb{N}_0.$$

Proof. By (2.6), the inequality (2.2) and [28, Theorem 1.6] the result follows. \square

Next, we recall the concept of discrete (α, ϱ) -resolvent sequence defined in [7, Section 4, Definition 4.4]. Also, useful results related with this definition are given.

Definition 2.5. *Let $\varrho, \alpha > 0$ be given and A be a closed linear operator with domain $D(A)$ defined on a Banach space X . An operator-valued sequence $\{S_{\alpha, \varrho}(n)\}_{n \in \mathbb{N}_0} \subset \mathcal{B}(X)$ is called a discrete (α, ϱ) -resolvent sequence generated by A if it satisfies the following conditions:*

- (i) $S_{\alpha, \varrho}(n)x \in D(A)$ for all $x \in X$ and $S_{\alpha, \varrho}(n)Ax = AS_{\alpha, \varrho}(n)x$ for each $n \in \mathbb{N}_0$ and $x \in D(A)$;
- (ii) $S_{\alpha, \varrho}(n)x = k^{\varrho}(n)x + A(k^{\alpha} * S_{\alpha, \varrho})(n)x$ for all $n \in \mathbb{N}_0$ and each $x \in X$.

We finish this section recalling the notion of (N, λ) -periodic sequences and their main properties. The notion of (N, λ) -periodic sequences was introduced in [6] as a discrete counterpart of the concept of (ω, c) -periodic functions defined in [10].

Definition 2.6 ([6]). *A vector-valued function $f : \mathbb{Z} \rightarrow X$ is called (N, λ) -periodic discrete function (or (N, λ) -periodic sequence) if there exist $N \in \mathbb{N}$ and $\lambda \in \mathbb{C} \setminus \{0\}$ such that $f(n + N) = \lambda f(n)$ for all $n \in \mathbb{Z}$. N is called the λ -period of f . The collection of those sequences with the same λ -period N will be denoted by $\mathbb{P}_{N\lambda}(\mathbb{Z}, X)$.*

The following result is central for the theory.

Proposition 2.7 ([6]). *A function f is (N, λ) -periodic discrete function if and only if there exists $u \in \mathbb{P}_N(\mathbb{Z}, X)$ such that*

$$(2.8) \quad f(n) = \lambda^{\wedge}(n)u(n), \quad \text{for all } n \in \mathbb{Z},$$

where $\lambda^{\wedge}(n) := \lambda^{n/N}$.

The vector-valued space of sequences $\mathbb{P}_{N\lambda}(\mathbb{Z}, X)$ is a Banach space with the norm

$$(2.9) \quad \|f\|_{N\lambda} := \max_{n \in [0, N]} \|\lambda^{(-n)} f(n)\|.$$

3. THE DISCRETE SCALED WRIGHT FUNCTION AND SUMMABILITY OF RESOLVENT SEQUENCES

In [7], the authors introduced a discrete version of the Lévy α -stable distribution which can be defined as

$$(3.1) \quad l_\alpha(n, j) = \sum_{i=0}^j \binom{j}{i} (-1)^i k^{-\alpha i}(n), \quad 0 < \alpha < 1, \quad n, j \in \mathbb{N}_0.$$

The sequence l_α is a probability density function in n , which means that

$$(3.2) \quad 0 \leq l_\alpha(n, j) \quad \text{and} \quad \sum_{n=0}^{\infty} l_\alpha(n, j) = 1.$$

This representation of the discrete Lévy function allowed to establish a subordination principle which relates a discrete (α, ϱ) -resolvent sequence with a C -semigroup generated by a given closed linear operator A defined on a Banach space X (see [7]). The following result is a consequence of this fact.

Theorem 3.1. *Let $0 < \alpha \leq \varrho \leq 1$ be given. Let A be a closed and linear operator defined on a Banach space X such that $1 \in \rho(A)$. Then the family*

$$(3.3) \quad S_{\alpha, \varrho}(n)x = \sum_{j=0}^{\infty} (k^{\varrho-\alpha} * l_\alpha(\cdot, j))(n) (I - A)^{-(j+1)} x, \quad n \in \mathbb{N}_0, \quad x \in X,$$

is a discrete (α, ϱ) -resolvent sequence generated by A .

Proof. By hypothesis, $C := (I - A)^{-1}$ exists and the operator $\{\mathcal{T}(n)\}_{n \in \mathbb{N}_0}$ given by $\mathcal{T}(n) = (I - A)^{-(n+1)}$ is bounded on X . On the other hand,

$$\mathcal{T}(n) = (I - A)^{-(n+1)} = C^{-(n-1)} \mathcal{T}(1)^n.$$

Hence, the operator $\{\mathcal{T}(n)\}_{n \in \mathbb{N}_0}$ is a the discrete C -semigroup (see [7]). Thus, the result follows from Theorem 4.5 of [7]. \square

The concept of scaled Wright function in the continuous case was introduced by Abadias and Miana in [5]. Motivated by the above theorem, we propose in this paper the following definition.

Definition 3.2. *Let $0 < \alpha < 1$ and $0 \leq \beta$ be given. For $n \in \mathbb{N}_0$ the discrete scaled Wright function $\varphi_{\alpha, \beta}$ is defined by*

$$(3.4) \quad \varphi_{\alpha, \beta}(n, j) := \sum_{i=0}^j \binom{j}{i} (-1)^i k^{\beta - \alpha i}(n).$$

Some properties of the discrete scaled Wright function can be deduced from those properties of the Lévy α -stable distribution. They are stated in the following remark.

Remark 3.3. (i) Note that, by (2.3) and (3.1)

$$\varphi_{\alpha,\beta}(n, j) = (k^\beta * l_\alpha(\cdot, j))(n)$$

and, in particular, $\varphi_{\alpha,0}(n, j) = l_\alpha(n, j)$.

(ii) Since the sequence l_α is non-negative, then the discrete scaled Wright function $\varphi_{\alpha,\beta}$ is non-negative.

(iii) The formula (3.3) can be written as

$$(3.5) \quad S_{\alpha,\varrho}(n) := \sum_{j=0}^{\infty} \varphi_{\alpha,\varrho-\alpha}(n, j) \mathcal{T}(j).$$

(iv) By Proposition 4.6 of [7], we have

$$\sum_{j=0}^{\infty} \varphi_{\alpha,\beta}(n, j) (1 + \lambda)^{-(j+1)} = \mathcal{E}_{\alpha,\alpha+\beta}(-\lambda, n), \quad n \in \mathbb{N}, \quad |\lambda| < 1.$$

(v) Let $A = \omega \in \mathbb{C}$ and $|\omega| < 1$. In this case, the discrete C -semigroup generated by A is given by $\mathcal{T}(n) = (1 - \omega)^{-(n+1)}$. Then, by Theorem 3.1, A generates discrete (α, α) -resolvent and $(\alpha, 1)$ -resolvent sequences given by

$$(3.6) \quad S_{\alpha,\alpha}(n) = \sum_{j=0}^{\infty} \varphi_{\alpha,0}(n, j) \mathcal{T}(j) = \mathcal{E}_{\alpha,\alpha}(\omega, n), \quad n \in \mathbb{N}_0,$$

and

$$(3.7) \quad S_{\alpha,1}(n) = \sum_{j=0}^{\infty} \varphi_{\alpha,1-\alpha}(n, j) \mathcal{T}(j) = \mathcal{E}_{\alpha,1}(\omega, n), \quad n \in \mathbb{N}_0,$$

where we have used (iii) and (iv).

We recall that an operator-valued sequence $\{S(n)\}_{n \in \mathbb{N}_0} \in \mathcal{B}(X)$ is said to be *summable* if

$$\|S\|_1 := \sum_{n=0}^{\infty} \|S(n)\| < \infty.$$

We finish this section with a useful result which is a direct consequence of the above considerations.

Theorem 3.4. Let A be a closed linear operator and suppose that $1 \in \rho(A)$ and

$$(3.8) \quad \|(I - A)^{-1}\| < 1.$$

Then A generates a summable discrete (α, α) -resolvent sequence $\{S_{\alpha,\alpha}(n)\}_{n \in \mathbb{N}_0}$.

Proof. Since $1 \in \rho(A)$, then by Theorem 3.1 the family

$$S_{\alpha,\alpha}(n)x = \sum_{j=0}^{\infty} \varphi_{\alpha,0}(n,j)(I - A)^{-(j+1)}x, \quad n \in \mathbb{N}_0, \quad x \in X,$$

is a discrete (α, α) -resolvent sequence generated by A . We will prove that it is summable. Indeed, since $0 \leq \varphi_{\alpha,0}(n,j) \leq 1$ for $j \in \mathbb{N}_0$, then

$$\begin{aligned} \sum_{n=0}^{\infty} \|S_{\alpha,\alpha}(n)\| &\leq \sum_{n=0}^{\infty} \sum_{j=0}^{\infty} \varphi_{\alpha,0}(n,j) \|(I - A)^{-(j+1)}\| \leq \sum_{j=0}^{\infty} \|(I - A)^{-(j+1)}\| \\ &\leq \|(I - A)^{-1}\| \sum_{j=0}^{\infty} \|(I - A)^{-1}\|^j < \infty. \end{aligned}$$

□

The following example provide concrete conditions on A under which the condition (3.8) holds.

Example 3.5. Let A be the generator of a C_0 -semigroup strictly contractive. For instance, on $X := L^1(\mathbb{R})$ we define

$$(T(t)f)(s) = \begin{cases} \beta f(t+s) & \text{if } s \in [-t, 0], \\ f(t+s) & \text{otherwise,} \end{cases}$$

where $0 < \beta < 1$ is arbitrary. Then $T(t)$ is a C_0 -semigroup and $\|T(t)\| = \beta < 1$ (since $\|T(t)\mathbb{1}_{[0,t]}\| = \beta\|\mathbb{1}_{[0,t]}\|$).

We deduce that $1 \in \rho(A)$ and $\|(I - A)^{-1}\| < 1$. Indeed,

$$\|(I - A)^{-1}\| = \left\| \int_0^{\infty} e^{-t} T(t) dt \right\| \leq \int_0^{\infty} e^{-t} \|T(t)\| dt < \beta < 1.$$

The last part of the earlier example shows the following result.

Corollary 3.6. Let A be the generator of a C_0 -semigroup strictly contractive, then $1 \in \rho(A)$ and $\|(I - A)^{-1}\| < 1$.

4. (N, λ) -PERIODIC SOLUTIONS FOR FRACTIONAL DIFFERENCE EQUATIONS ON \mathbb{Z}

In this section, we study regularity of solutions to the linear fractional difference equation

$$\Delta^\alpha u(n) = Au(n+1) + g(n), \quad n \in \mathbb{Z},$$

and the nonlinear fractional equation

$$\Delta^\alpha u(n) = Au(n+1) + f(n, u(n)), \quad n \in \mathbb{Z}.$$

in $\mathbb{P}_{N\lambda}(\mathbb{Z}, X)$, where A be a closed linear operator with domain $D(A)$ defined on X .

4.1. The linear case.

Let $0 < \alpha \leq 1$ and A be a closed linear operator with domain $D(A)$ defined on a Banach space X . We consider the linear fractional difference equation

$$(4.1) \quad \Delta^\alpha u(n) = Au(n+1) + g(n), \quad n \in \mathbb{Z}.$$

We recall from [1, Definition 4.1] that a sequence $u \in \ell^1(\mathbb{Z}, X)$ is called a *strong solution* for equation (4.1) if $u(n) \in D(A)$ for all $n \in \mathbb{Z}$ and u satisfies (4.1).

Definition 4.1 ([1]). *Let A be the generator of a discrete (α, α) -resolvent family $\{S_{\alpha, \alpha}(n)\}_{n \in \mathbb{N}_0}$ and $g : \mathbb{Z} \rightarrow X$. The sequence*

$$(4.2) \quad u(n) = \sum_{j=-\infty}^{n-1} S_{\alpha, \alpha}(n-1-j)g(j), \quad n \in \mathbb{Z}.$$

is called a mild solution for equation (4.1) if $m \rightarrow S_{\alpha, \alpha}(m)g(n-m)$ is summable on \mathbb{N}_0 for each $n \in \mathbb{Z}$.

Note that if $g \in \ell^1(\mathbb{Z}, D(A))$ then each mild solution is a strong one, see [1, Theorem 4.2].

In the following theorem, we establish the existence of (N, λ) -periodic mild solutions for equation (4.1).

Theorem 4.2. *Let $0 < \alpha \leq 1$. Assume that A be a closed linear operator defined on a Banach space X , $1 \in \rho(A)$ and*

$$\|(I - A)^{-1}\| < 1.$$

If $g \in \mathbb{P}_{N\lambda}(\mathbb{Z}, X)$, then there is an (N, λ) -periodic mild solution of (4.1) given by the sequence

$$(4.3) \quad u(n) := \sum_{j=-\infty}^{n-1} S_{\alpha, \alpha}(n-1-j)g(j), \quad n \in \mathbb{Z},$$

where $\{S_{\alpha, \alpha}(n)\}_{n \in \mathbb{N}_0}$ is discrete (α, α) -resolvent sequence defined in (3.3).

Proof. By Theorem 3.4, A generates a summable discrete (α, α) -resolvent sequence $\{S_{\alpha, \alpha}(n)\}_{n \in \mathbb{N}}$ given by

$$S_{\alpha, \alpha}(n)x = \sum_{j=0}^{\infty} \varphi_{\alpha, 0}(n, j)(I - A)^{-(j+1)}x, \quad n \in \mathbb{N}_0, \quad x \in X.$$

Since g is bounded and $\{S_{\alpha, \alpha}(n)\}_{n \in \mathbb{N}_0}$ is summable, it follows that the sequence u is a mild solution of (4.1).

It remains to prove that $u \in \mathbb{P}_{N\lambda}(\mathbb{Z}, X)$.

Indeed,

$$\begin{aligned} u(n+N) &= \sum_{j=-\infty}^{n+N-1} S_{\alpha,\alpha}(n+N-1-j)g(j) = \sum_{p=-\infty}^{n-1} S_{\alpha,\alpha}(n-1-p)g(p+N) \\ &= \lambda \sum_{p=-\infty}^{n-1} S_{\alpha,\alpha}(n-1-p)g(p) = \lambda u(n), \end{aligned}$$

getting that $u \in \mathbb{P}_{N\lambda}(\mathbb{Z}, X)$.

□

4.2. The semilinear case.

In this subsection, we consider the following fractional difference equation

$$(4.4) \quad \Delta^\alpha u(n) = Au(n+1) + f(n, u(n)), \quad n \in \mathbb{Z},$$

where $0 < \alpha \leq 1$, A satisfies the hypotheses in Theorem 3.4 and f satisfies suitable conditions.

Inspired in the solution of the linear case, we give the following definition of mild solution for the semilinear case.

Definition 4.3. *Let A be the generator of a discrete (α, α) -resolvent family $\{S_{\alpha,\alpha}(n)\}_{n \in \mathbb{N}_0}$ and $f : \mathbb{Z} \times X \rightarrow X$. We say that a sequence $u : \mathbb{Z} \rightarrow X$ is a (N, λ) -periodic mild solution of (4.4) if $u \in \mathbb{P}_{N\lambda}(\mathbb{Z}, X)$, satisfies*

$$(4.5) \quad u(n) = \sum_{j=-\infty}^{n-1} S_{\alpha,\alpha}(n-1-j)f(j, u(j)), \quad n \in \mathbb{Z},$$

where $m \rightarrow S_{\alpha,\alpha}(m)f(n-m, x)$ is summable on \mathbb{N}_0 for each $n \in \mathbb{Z}$.

Let $f : \mathbb{Z} \times X \rightarrow X$, $\phi \in \mathbb{P}_{N\lambda}(\mathbb{Z}, X)$ and denote by $\mathcal{N}(\phi)(\cdot) := f(\cdot, \phi(\cdot))$ the Nemytskii discrete composition operator.

To prove the main theorem we will need to recall the following.

Theorem 4.4 ([6]). *Let $f : \mathbb{Z} \times X \rightarrow X$. Then the following assertions are equivalent:*

- (i) *For every $\phi \in \mathbb{P}_{N\lambda}(\mathbb{Z}, X)$ we have that $\mathcal{N}(\phi)$ is (N, λ) -periodic discrete.*
- (ii) *f is N -periodic in the first variable and homogeneous in the second variable, that is $f(n+N, \lambda x) = \lambda f(n, x)$ for all $(n, x) \in \mathbb{Z} \times X$.*

Let $\mathbb{D} := \{\lambda \in \mathbb{C} : |\lambda| < 1\}$. The following is our main result.

Theorem 4.5. *Let $f : \mathbb{Z} \times X \rightarrow X$ be given and let A be a closed linear operator defined on a Banach space X such that $1 \in \rho(A)$ and*

$$(4.6) \quad r_A := \|(I - A)^{-1}\| < 1.$$

Assume the following conditions:

H_1 . There exists $(N, \lambda) \in \mathbb{N} \times (\mathbb{C} \setminus \mathbb{D})$ such that $f(n + N, \lambda x) = \lambda f(n, x)$ for all $(n, x) \in \mathbb{Z} \times X$.

H_2 . There exists a constant $L > 0$ such that

$$\|f(n, x) - f(n, y)\| \leq L\|x - y\|,$$

for all $x, y \in X$ and all $n \in \mathbb{Z}$.

H_3 . The constant L in H_2 is such that

$$L < \left(1 - \frac{1}{|\lambda|^{1/N}}\right)^\alpha + \left(\frac{1}{r_A} - 1\right).$$

Then, equation (4.4) has a unique (N, λ) -periodic mild solution.

Proof. First, let us define the operator $G : \mathbb{P}_{N\lambda}(\mathbb{Z}, X) \rightarrow \mathbb{P}_{N\lambda}(\mathbb{Z}, X)$ by

$$G(u)(n) := \sum_{j=-\infty}^{n-1} S_{\alpha, \alpha}(n-1-j) f(j, u(j)).$$

Let $u \in \mathbb{P}_{N\lambda}(\mathbb{Z}, X)$ and $g(n) := f(n, u(n))$. By H_1 and Theorem 4.4 we get that $g \in \mathbb{P}_{N, \lambda}(\mathbb{Z}, X)$. As in the linear case, we can see that $G(u) \in \mathbb{P}_{N\lambda}(\mathbb{Z}, X)$. It follows that G is well defined. Now, for $u, v \in \mathbb{P}_{N\lambda}(\mathbb{Z}, X)$,

$$\|G(u) - G(v)\|_{N\lambda} = \max_{n \in [0, N]} \|\lambda^{-(n-1)} \sum_{j=-\infty}^{n-1} S_{\alpha, \alpha}(n-1-j) [f(j, u(j)) - f(j, v(j))]\|$$

where we have by H_2 that

$$\begin{aligned} & \|\lambda^{-(n-1)} \sum_{j=-\infty}^{n-1} S_{\alpha, \alpha}(n-1-j) [f(j, u(j)) - f(j, v(j))]\| \\ &= \left\| \sum_{j=-\infty}^{n-1} \lambda^{-(n-1-j)} S_{\alpha, \alpha}(n-1-j) \lambda^{-(j)} [f(j, u(j)) - f(j, v(j))]\right\| \\ &< \sum_{j=-\infty}^{n-1} |\lambda|^{-(n-1-j)} |S_{\alpha, \alpha}(n-1-j)| |\lambda|^{-j} \| [f(j, u(j)) - f(j, v(j))]\| \\ &< L \sum_{j=-\infty}^{n-1} |\lambda|^{-(n-1-j)} |S_{\alpha, \alpha}(n-1-j)| \|\lambda^{-(j)} [u(j) - v(j)]\| \\ &< \|u - v\|_{N\lambda} L \sum_{k=0}^{\infty} \|S_{\alpha, \alpha}^{\sim}(k)\|, \end{aligned}$$

where $S_{\alpha,\alpha}^{\sim}(n) = \lambda^{(-n)}S_{\alpha,\alpha}(n)$. Then,

$$\begin{aligned} \|G(u) - G(v)\|_{N\lambda} &= \max_{n \in [0, N]} \|\lambda^{(-n)} \sum_{j=-\infty}^n S_{\alpha,\alpha}(n-1-j) [f(j, u(j)) - f(j, v(j))]\| \\ &\leq L\|u - v\|_{N\lambda} \|S_{\alpha,\alpha}^{\sim}\|_1 < \|u - v\|_{N\lambda}, \end{aligned}$$

where by Theorem 3.1, Remark 3.3 (iv) and (2.7) we have

$$\begin{aligned} \|S_{\alpha,\alpha}^{\sim}\|_1 &\leq \sum_{n=0}^{\infty} |\lambda|^{-n/N} \sum_{j=0}^{\infty} \varphi_{\alpha,0}(n, j) r_A^{j+1} = \sum_{n=0}^{\infty} |\lambda|^{-n/N} \mathcal{E}_{\alpha,\alpha} \left(1 - \frac{1}{r_A}, n\right) \\ &= \frac{|\lambda|^{\alpha/N}}{(|\lambda|^{1/N} - 1)^{\alpha} - (1 - \frac{1}{r_A})|\lambda|^{\alpha/N}} = \frac{1}{\left(1 - \frac{1}{|\lambda|^{1/N}}\right)^{\alpha} + \left(\frac{1}{r_A} - 1\right)}. \end{aligned}$$

Therefore, the conclusion follows from H_3 . For the above, it follows that there exists a unique function $u \in \mathbb{P}_{N\lambda}(\mathbb{Z}, X)$ such that $G u = u$. Hence u is the unique (N, λ) -periodic mild solution of equation (4.4). \square

Remark 4.6. Regarding condition H_3 we observe that is enough to have the weaker condition $L\|S_{\alpha,\alpha}^{\sim}\|_1 < 1$ where $S_{\alpha,\alpha}^{\sim}(n) := \lambda^{(-n)}S_{\alpha,\alpha}(n)$ and $\{S_{\alpha,\alpha}(n)\}_{n \in \mathbb{N}_0}$ is the (α, α) -resolvent sequence generated by A .

We finally finish with an application of the main result presented in this paper.

Example 4.7. Let $0 < \alpha < 1$ and $|\lambda| \geq 1$. We consider the following fractional difference-differential equation in $X = L^2(0, 1)$

$$(4.7) \quad \begin{cases} \Delta^{\alpha} u(n, x) = \frac{\partial^2}{\partial x^2} u(n+1, x) + g(n, x) \cos(h(n, x)u(n, x)), & n \in \mathbb{Z}, x \in (0, 1), \\ u(n, 0) = u(n, 1) = 0, \end{cases}$$

where $g \in \mathbb{P}_{N\lambda}(\mathbb{Z}, L^2(0, 1))$, $h \in \mathbb{P}_{N\frac{1}{\lambda}}(\mathbb{Z}, L^2(0, 1))$ and

$$(4.8) \quad \max_{n \in [0, N]} \|g(n)h(n)\|_{L^2} < (1 - |\lambda|^{-1/N})^{\alpha} + \left(\left(\sum_{m=1}^{\infty} \frac{1}{(1 + (m\pi)^2)^2} \right)^{-1/2} - 1 \right).$$

We define

$$\begin{aligned} D(A) &= \{f \in L^2(0, 1) : f'' \in L^2(0, 1), f(0) = f(1) = 0\}, \\ Af &= f'', \quad \forall f \in D(A). \end{aligned}$$

Then (4.7) can be written in the abstract setting (4.4). It is well known that A is the generator of an analytic semigroup $\{T(t)\}_{t \geq 0}$ on $L^2(0, 1)$ (see [19]) which is given by

$$T(t)f = \sum_{j=0}^{\infty} e^{-j^2\pi^2 t} \langle f, \mathbf{e}_j \rangle \mathbf{e}_j,$$

where $\{\mathbf{e}_j\}$ is the standard basis in $L^2(0, 1)$. Moreover, we can represent the generator A as

$$Af = - \sum_{m=1}^{\infty} (m\pi)^2 \langle f, \mathbf{e}_m \rangle \mathbf{e}_m, \quad f \in D(A).$$

Then, for each $f \in L^2(0, 1)$ we have $1 \in \rho(A)$ and

$$\|(I - A)^{-1}f\|_{L^2}^2 = \sum_{m=1}^{\infty} \frac{1}{(1 + (m\pi)^2)^2} |\langle f, \mathbf{e}_m \rangle|^2.$$

Note that

$$r_A := \sup_{\|f\|_{L^2}=1} \|(I - A)^{-1}f\|_{L^2} = \left(\sum_{m=1}^{\infty} \frac{1}{(1 + (m\pi)^2)^2} \right)^{1/2} \leq \left(\frac{1}{\pi^4} \sum_{m=1}^{\infty} \frac{1}{m^4} \right)^{1/2} = \frac{1}{\sqrt{90}} < 1,$$

where we have used the formula [29, Pag. 651] in the last equality. Then, the condition (4.6) is satisfied. Now, we shall verify all the hypotheses in Theorem 4.5. Indeed, the sequence $f(n, \xi) := g(n) \cos(h(n)\xi)$, $\xi \in L^2(0, 1)$, satisfies:

$$\begin{aligned} f(n + N, \lambda\xi) &= g(n + N) \cos(h(n + N)\lambda\xi) = \lambda g(n) \cos\left(\frac{1}{\lambda}h(n)\lambda\xi\right) \\ &= \lambda g(n) \cos(h(n)\xi) = \lambda f(n, \xi), \end{aligned}$$

and

$$\|f(n, \xi) - f(n, \psi)\|_{L^2} \leq \|g(n)h(n)\|_{L^2} \|\xi - \psi\|_{L^2} \leq L \|\xi - \psi\|_{L^2},$$

where

$$L := \max_{n \in [0, N]} \|g(n)h(n)\|_{L^2}.$$

From equation (4.8) and the fact that $r_A < 1$, we obtain that

$$L < \left(1 - \frac{1}{|\lambda|^{1/N}}\right)^\alpha + \left(\frac{1}{r_A} - 1\right),$$

satisfying H_3 . Thus, we have checked all the hypotheses of Theorem 4.5. Hence equation (4.7) has a unique (N, λ) -periodic mild solution.

Finally, observe that in case $|\lambda| = 1$, we have that

$$L := \max_{n \in [0, N]} \|g(n)h(n)\|_{L^2} < \frac{1}{r_A} - 1.$$

and therefore condition H_3 independent of α . This happens precisely in the standard cases of discrete periodic, discrete anti-periodic and discrete Bloch periodic functions.

ACKNOWLEDGMENTS

We thank Professor William Ramirez for his hospitality and the Universidad de la Costa (CUC) for allowing the use of its facilities.

REFERENCES

- [1] L. Abadias, C. Lizama. *Almost automorphic mild solutions to fractional partial difference-differential equations*, Appl. Anal. 95 (6) (2016), 1347–1369.
- [2] L. Abadias, C. Lizama, P. J. Miana, M. P. Velasco. *Cesàro sums and algebra homomorphisms of bounded operators*, Israel J. Math. 216 (1) (2016), 471–505.
- [3] T. Abdeljawad, S. Banerjee, G. C. Wu. *Discrete tempered fractional calculus for new chaotic systems with short memory and image encryption*. Optik. 218 (2020), 163698.
- [4] M. Agaoglou, M. Fečkan, A. Panagiotidou. *Existence and uniqueness of (ω, c) -periodic solutions of semilinear evolution equations*, Int. J. Dyn. Sys. and Diff. Equ. 10 (2) (2020), 149–166.
- [5] L. Abadias, P. Miana. *A subordination principle on Wright functions and regularized resolvent families*, J. Funct. Spaces, (2015), 2314–889.
- [6] E. Alvarez, S. Díaz, C. Lizama. *On the existence and uniqueness of (N, λ) -periodic solutions to a class of Volterra difference equations*, Adv. Differ. Equ. 105 (2019).
- [7] E. Alvarez, S. Díaz, C. Lizama. *C-Semigroups, subordination principle and the Lévy α -stable distribution on discrete time*, Commun. Contemp. Math., to appear. <https://doi.org/10.1142/S0219199720500637>
- [8] E. Alvarez, S. Castillo, M. Pinto. *(ω, c) -asymptotically periodic functions, first-order Cauchy problem, and Lasota-Ważewska model with unbounded oscillating production of red cells*. Math. Meth. Appl. Sci. 43 (1) (2020), 305–319.
- [9] E. Alvarez, S. Castillo, M. Pinto. *(ω, c) -pseudo periodic functions, first order Cauchy problem and Lasota-Ważewska model with ergodic and unbounded oscillating production of red cells*. Bound. Value Probl. 2019, 106 (2019). <https://doi.org/10.1186/s13661-019-1217-x>
- [10] E. Alvarez, A. Gómez, M. Pinto. *(ω, c) -periodic functions and mild solutions to abstract fractional integro-differential equations*. Electr. J. Qual. Th. Diff. Equ. (16) (2018), 1–8.
- [11] C. Goodrich, C. Lizama. *A transference principle for nonlocal operators using a convolutional approach: fractional monotonicity and convexity*, Isr. J. Math. 236 (2020), 533–589.
- [12] J. W. He, C. Lizama, Y. Zhou. *The Cauchy problem for discrete-time fractional evolution equations*. J. Comput. Appl. Math., 370 (2020), 112683.
- [13] L. L. Huang, G. C. Wu, D. Baleanu, H.Y. Wang. *Discrete fractional calculus for interval-valued systems*. Fuzzy Sets and Systems. 404 (2021), 141–158.
- [14] L. L. Huang, J. H. Park, G. C. Wu, Z. W. Mo. *Variable-order fractional discrete-time recurrent neural networks*. J. Comp. Appl. Math. 370 (2020), 112633
- [15] M. Khalladi, M. Kostić, A. Rahmani, D. Velinov. *(ω, c) -Almost periodic type functions and applications*. 2020, hal.archives-ouvertes.fr
- [16] M. Khalladi, A. Rahmani. *(ω, c) -Pseudo almost periodic distributions*. Nonautonomous Dynamical Systems, 7 (1) (2020).
- [17] M. Kostić, V.E. Fedorov. *Asymptotically (ω, c) -almost periodic type solutions of abstract degenerate non-scalar Volterra equations*. Chelyabinsk Physical and Mathematical Journal. 5 (4) (2020), 415–427.
- [18] M. Kostić. *Multidimensional (ω, c) - almost periodic type functions and applications*. hal.archives-ouvertes.fr
- [19] A. Lunardi. *Analytic Semigroups and Optimal Regularity in Parabolic Problems*. PNLDE, Vol. 16, Birkhäuser Verlag, Basel, 1995.
- [20] K. Liu, J. Wang, D. O’Regan, M. Fečkan. *A new class of (ω, c) -periodic non-instantaneous impulsive differential equations*. Mediterr. J. Math. 17 (2020), 155.
- [21] C. Lizama. *The Poisson distribution, abstract fractional difference equations, and stability*, Proc. Amer. Math. Soc., 145 (9) (2017), 3809–3827.
- [22] C. Lizama. *ℓ_p -maximal regularity for fractional difference equations on UMD spaces*, Math. Nach., 288 (17/18) (2015), 2079–2092.
- [23] C. Lizama, M. Murillo. *Maximal regularity in ℓ_p spaces for discrete time fractional shifted equations*. J. Diff. Equations, 263 (6) (2017), 3175–3196.

- [24] C. Lizama, M.P. Velasco. *Weighted bounded solutions for a class of nonlinear fractional equations*. *Fract. Calculus Appl. Anal.*, 19 (4) (2016), 1010–1030.
- [25] C. Lizama, R. Ponce. *Solutions of abstract integro-differential equations via Poisson transformation*, *Math Meth Appl Sci.* 44(3) (2021), 2495–2505.
- [26] G. Mophou, G. N’Guérékata. *An existence result of (ω, c) -periodic mild solutions to some fractional differential equation*. hal.archives-ouvertes.fr
- [27] D. Mozyrska, M. Wyrwas. *The Z-transform method and delta type fractional difference operators*, *Disc. Dyn. Nat. Soc.*, 2015, Art. ID 852734, 12,1026-0226, doi:10.1155/2015/852734.
- [28] I. Podlubny. *Fractional Differential Equations*, Vol. 198, Academic Press, 1999, 340.
- [29] A. P. Prudnikov, A. Y. Brychkov, and O. I. Marichev. *Integrals and series. Vol. 1. Elementary functions*, Gordon and Breach Science Publishers, New York, 1986.
- [30] J. Wang, L. Ren, Y. Zhou. *(ω, c) -periodic solutions for time varying impulsive differential equations*, *Adv. Differ. Equ.* (2019), 259.
- [31] G. C. Wu, D. Baleanu, Y. R. Bai. *Discrete fractional masks and their applications to image enhancement*. *Appl. in Eng., Life and Social Sc., Part B* (2019), 261–270.
- [32] G. C. Wu, D. Baleanu, W. H. Luo. *Lyapunov functions for Riemann-Liouville-like fractional difference equations*. *Appl. Math. Comp.* 314 (2017), 228–236.
- [33] A. Zygmund. *Trigonometric Series*, 2nd ed. Vol. I Cambridge University Press, New York, 1959.

UNIVERSIDAD DEL NORTE, DEPARTAMENTO DE MATEMÁTICAS Y ESTADÍSTICA, BARRANQUILLA, COLOMBIA.

Email address: ealvareze@uninorte.edu.co

UNIVERSIDAD DEL NORTE, DEPARTAMENTO DE MATEMÁTICAS Y ESTADÍSTICA, BARRANQUILLA, COLOMBIA.

Email address: stivend@uninorte.edu.co

UNIVERSIDAD DE SANTIAGO DE CHILE (USACH), FACULTAD DE CIENCIA, DEPARTAMENTO DE MATEMÁTICA Y CIENCIA DE LA COMPUTACIÓN, LAS SOPHORAS 173, ESTACIÓN CENTRAL, SANTIAGO, CHILE.

Email address: carlos.lizama@usach.cl