

MAXIMAL REGULARITY FOR DEGENERATE DIFFERENTIAL EQUATIONS WITH INFINITE DELAY IN PERIODIC VECTOR-VALUED FUNCTION SPACES.

CARLOS LIZAMA AND RODRIGO PONCE

ABSTRACT. Let A and M be closed linear operators defined on a complex Banach space X and $a \in L^1(\mathbb{R}_+)$ be an scalar kernel. We use operator-valued Fourier multipliers techniques to obtain necessary and sufficient conditions to guarantee the existence and uniqueness of periodic solutions to the equation

$$\frac{d}{dt}(Mu(t)) = \alpha Au(t) + \int_{-\infty}^t a(t-s)Au(s)ds + f(t), \quad t > 0,$$

with initial condition $Mu(0) = Mu(2\pi)$, solely in terms of spectral properties of the data. Our results are obtained in the scales of periodic Besov, Triebel-Lizorkin and Lebesgue vector-valued function spaces.

1. INTRODUCTION

In this paper, we study maximal regularity in Lebesgue, Besov and Triebel-Lizorkin vector-valued function spaces for the following class of differential equation with infinite delay

$$(1.1) \quad \frac{d}{dt}(Mu(t)) = \alpha Au(t) + \int_{-\infty}^t a(t-s)Au(s)ds + f(t), \quad 0 \leq t \leq 2\pi,$$

where $(A, D(A))$ and $(M, D(M))$ are (unbounded) closed linear operators defined on a Banach space X , with $D(A) \subseteq D(M)$, $a \in L^1(\mathbb{R}_+)$ an scalar-valued kernel, $\alpha \in \mathbb{R} \setminus \{0\}$ and f an X -valued function defined on $[0, 2\pi]$.

The model (1.1) corresponds to problems related with viscoelastic materials; that is, materials whose stresses at any instant depend on the complete history of strains that the material has undergone (see [21]) or heat conduction with memory. For more details, see, for instance, [14], [15] and [23].

The recent linear theory of maximal regularity is not only important on its own but it is also the indispensable basis for the theory of nonlinear evolution equations, see e.g. [1], [13], [23] and references therein. In case $M = I$ (the identity in X) and $a \equiv 0$, equation (1.1) with periodic initial conditions have been studied by Arendt-Bu, Bu-Kim

2000 *Mathematics Subject Classification.* 34G10, 34K13, 45N05, 45D05.

Key words and phrases. Differential equations with delay; Operator-valued Fourier multipliers; R -boundedness; UMD -spaces; Besov vector valued spaces; Triebel-Lizorkin vector valued spaces; Lebesgue vector valued spaces.

The first author is partially supported by FONDECYT Grant #1100485.

and characterizations of the maximal regularity in Lebesgue, Besov and Triebel-Lizorkin vector-valued function spaces were obtained using the resolvent set of A . See [3], [4] and [7].

On the other hand, characterizations of maximal regularity for equation (1.1) in case $M = I$ and $a \in L^1(\mathbb{R})$ have been obtained by Keyantuo-Lizama [18] in Lebesgue and Besov vector-valued function spaces and by Bu-Fang [8] in Triebel-Lizorkin vector valued spaces. We note that periodic solutions have been also studied by other authors, for example in [11] using topological methods.

Characterizations of maximal regularity in these vector-valued function spaces, for the degenerate abstract equation (1.1) with periodic initial conditions

$$(1.2) \quad Mu(0) = Mu(2\pi),$$

and $a \equiv 0$ has been studied recently by the authors [22]. The method used in [22] is based on the results given by Arendt-Bu ([3, 4]) and Bu-Kim ([7]), for operator-valued Fourier multipliers in Lebesgue, Besov and Triebel-Lizorkin vector-valued function spaces. It is worthwhile to mention that this method enables to improve and extend results on degenerate abstract equations obtained previously in the literature. Compare e.g. Barbu-Favini [5] and [22].

In this article, we apply the same method to obtain characterizations of maximal regularity for (1.1) in the above mentioned vector-valued function spaces. The advantage of our approach is clear. We recover, as special cases, the results in [3], [4], [7], [8], [18] and [22]. It is remarkable that in the characterizations that we will obtain, no conditions on the commutativity of operators A and M , or in the existence of bounded inverse of A or M , are needed. Also, in the case of periodic Besov and Triebel-Lizorkin function spaces, no geometrical assumption on the underlying Banach space X is needed.

The plan of the paper is the following: After some preliminaries in the second section, assuming that X is an UMD space, we characterize in Section 3 the uniqueness and existence of a strong L^p -solution for the problem (1.1)-(1.2) solely in terms of a property of R -boundedness for the sequence of operators $ikM(ikM - (\alpha + \tilde{a}(ik))A)^{-1}$. Here the tilde denotes Laplace transform of $a(t)$. In Section 4, we obtain a characterization in the context of Besov spaces. We notice that, as particular case of this characterization, a simply condition to guarantee the existence and uniqueness of solution in Hölder spaces $C^s((0, 2\pi); X)$, $0 < s < 1$, in general Banach spaces X , is obtained. Namely

Theorem 1.1. *Let $s > 0$ and $A : D(A) \subseteq X \rightarrow X$, $M : D(M) \subseteq X \rightarrow X$ be closed linear operators on a Banach space X . Suppose that $D(A) \subseteq D(M)$, the sequence $\{\tilde{a}(ik)\}_{k \in \mathbb{Z}}$ is 2-regular and $(ikM - (\alpha + \tilde{a}(ik))A)$ are closed operators for all $k \in \mathbb{Z}$ where $\alpha \neq 0$. Then, the following assertions are equivalent:*

(i) *For every $f \in C^s((0, 2\pi); X)$ there is a unique strong C^s -solution of (1.1) such that $Mu(0) = Mu(2\pi)$;*

(ii) *$(ikM - (\alpha + \tilde{a}(ik))A)^{-1}$ exists for all $k \in \mathbb{Z}$ and $\sup_{k \in \mathbb{Z}} \|(ikM - (\alpha + \tilde{a}(ik))A)^{-1}\| < \infty$.*

In Section 5 we give the corresponding characterization in case of the scale of Triebel-Lizorkin vector valued spaces. The difference with the scale of Besov vector valued spaces is only that we need more regularity of the sequence $\tilde{a}(ik)$. In Section 6, we apply our results in two concrete examples.

2. PRELIMINARIES

Given $1 \leq p < \infty$, we denote by $L_{2\pi}^p(\mathbb{R}, X)$ the space of all 2π -periodic Bochner measurable X -valued functions f , such that the restriction of f to $[0, 2\pi]$ is p -integrable. For a function $f \in L_{2\pi}^1(\mathbb{R}, X)$ we denote by $\hat{f}(k)$, the k -th Fourier coefficient of f :

$$\hat{f}(k) = \frac{1}{2\pi} \int_0^{2\pi} e^{-ikt} f(t) dt,$$

for all $k \in \mathbb{Z}$. We remark that the Fourier coefficients determine the function f , that is, $\hat{f}(k) = 0$ for all $k \in \mathbb{Z}$ if and only if $f(t) = 0$ a.e. Let X, Y be Banach spaces. We denote by $\mathcal{B}(X, Y)$ be the space of all bounded linear operators from X to Y . When $X = Y$, we write simply $\mathcal{B}(X)$. For a linear operator A on X , we denote domain by $D(A)$ and its resolvent set by $\rho(A)$. By $[D(A)]$ we denote the domain of A equipped with the graph norm.

We begin with some preliminaries about operator-valued Fourier multipliers. More information can be found in [4] in the periodic case and for the non-periodic case, see for instance, [25].

Definition 2.1. [3] *For $1 \leq p < \infty$, we say that a sequence $\{M_k\}_{k \in \mathbb{Z}} \subset \mathcal{B}(X, Y)$ is an L^p -multiplier if, for each $f \in L_{2\pi}^p(\mathbb{R}, X)$, there exists $u \in L_{2\pi}^p(\mathbb{R}, Y)$ such that*

$$\hat{u}(k) = M_k \hat{f}(k) \text{ for all } k \in \mathbb{Z}.$$

It follows from the uniqueness theorem of Fourier series that u is uniquely determined by f .

For $j \in \mathbb{N}$, denote by r_j the j -th Rademacher function on $[0, 1]$ i.e. $r_j(t) = \text{sgn}(\sin(2^j \pi t))$ and for $x \in X$, $r_j \otimes x$, denote the vector valued function $t \rightarrow r_j(t)x$.

Definition 2.2. *A family of operators $\mathcal{T} \subset \mathcal{B}(X, Y)$ is called R -bounded, if there is a constant $C_p > 0$ and $p \in [1, \infty)$ such that for each $N \in \mathbb{N}, T_j \in \mathcal{T}, x_j \in X, j = 1, \dots, N$ the inequality*

$$(2.1) \quad \left\| \sum_{j=1}^N r_j \otimes T_j x_j \right\|_{L^p((0,1);Y)} \leq C_p \left\| \sum_{j=1}^N r_j \otimes x_j \right\|_{L^p((0,1);X)}$$

is valid.

If (2.1) holds for some $p \in [1, \infty)$ then it holds for all $p \in [1, \infty)$. The smallest C_p in (2.1) is called R -bound of \mathcal{T} , we denote it by $R_p(\mathcal{T})$.

We remark that large classes of classical operators are R -bounded (cf. [16] and references therein). Hence, this assumption is not too restrictive for the applications that we consider in this article.

Remark 2.3.

Several properties of R -bounded families can be founded in the monograph of Denk-Hieber-Prüss [13]. For the reader's convenience, we summarize here from [13, Section 3] some results.

(a) If $\mathcal{T} \subset \mathcal{B}(X, Y)$ is R -bounded then it is uniformly bounded, with

$$\sup\{\|T\| : T \in \mathcal{T}\} \leq R_p(\mathcal{T}).$$

(b) The definition of R -boundedness is independent of $p \in [1, \infty)$.

(c) When X and Y are Hilbert spaces, $\mathcal{T} \subset \mathcal{B}(X, Y)$ is R -bounded if and only if \mathcal{T} is uniformly bounded.

(d) Let X, Y be Banach spaces and $\mathcal{T}, \mathcal{S} \subset \mathcal{B}(X, Y)$ be R -bounded. Then

$$\mathcal{T} + \mathcal{S} = \{T + S : T \in \mathcal{T}, S \in \mathcal{S}\}$$

is R -bounded as well, and $R_p(\mathcal{T} + \mathcal{S}) \leq R_p(\mathcal{T}) + R_p(\mathcal{S})$.

(e) Let X, Y, Z be Banach spaces, and $\mathcal{T} \subset \mathcal{B}(X, Y)$ and $\mathcal{S} \subset \mathcal{B}(Y, Z)$ be R -bounded. Then

$$\mathcal{ST} = \{ST : T \in \mathcal{T}, S \in \mathcal{S}\}$$

is R -bounded, and $R_p(\mathcal{ST}) \leq R_p(\mathcal{S})R_p(\mathcal{T})$.

(g) Let X, Y be Banach spaces and $\mathcal{T} \subset \mathcal{B}(X, Y)$ is R -bounded. If $\{\alpha_k\}_{k \in \mathbb{Z}}$ is a bounded sequence, then $\{\alpha_k T : T \in \mathcal{T}\}$ is R -bounded.

Proposition 2.4. [3] *Let X be a Banach space and $\{M_k\}_{k \in \mathbb{Z}}$ be an L^p -multiplier, where $1 \leq p < \infty$. Then, the set $\{M_k : k \in \mathbb{Z}\}$ is R -bounded.*

A Banach space X is said to be UMD , if the Hilbert transform is bounded on $L^p(\mathbb{R}, X)$ for some (and then all) $p \in (1, \infty)$. Here the Hilbert transform H of a function $f \in \mathcal{S}(\mathbb{R}, X)$, the Schwartz space of rapidly decreasing X -valued functions, is defined by

$$Hf := \frac{1}{\pi} PV\left(\frac{1}{t}\right) * f.$$

These spaces are also called \mathcal{HT} spaces. It is a well known that the set of Banach spaces of class \mathcal{HT} coincides with the class of UMD spaces. This has been shown by Bourgain [6] and Burkholder [9]. Some examples of UMD -spaces include the Hilbert spaces, Sobolev spaces $W_p^s(\Omega)$, $1 < p < \infty$, Lebesgue spaces $L^p(\Omega, \mu)$, $1 < p < \infty$, $L^p(\Omega, \mu; X)$, $1 < p < \infty$, when X is a UMD -space. Moreover, a UMD -space is reflexive and therefore, $L^1(\Omega, \mu)$, $L^\infty(\Omega, \mu)$ (in the case infinite dimensional) and $C^s([0, 2\pi]; X)$ are not UMD . More information on UMD spaces can be found in [6, 9] and [10].

We recall the following theorem, due to Arendt-Bu [3, Theorem 1.3], in the context of UMD -spaces.

Theorem 2.5. [3] *Let X, Y be UMD spaces and let $\{M_k\}_{k \in \mathbb{Z}} \subseteq \mathcal{B}(X, Y)$. If the sets $\{M_k\}_{k \in \mathbb{Z}}$ and $\{k(M_{k+1} - M_k)\}_{k \in \mathbb{Z}}$ are R -bounded, then $\{M_k\}_{k \in \mathbb{Z}}$ is an L^p -multiplier for $1 < p < \infty$.*

We will need the following Lemmas.

Lemma 2.6. [3] *Let $f, g \in L_{2\pi}^p(\mathbb{R}; X)$, where $1 \leq p < \infty$ and A is a closed operator on a Banach space X . Then, the following assertions are equivalent.*

(i) $f(t) \in D(A)$ and $Af(t) = g(t)$, a.e.

(ii) $\hat{f}(k) \in D(A)$ and $A\hat{f}(k) = \hat{g}(k)$, for all $k \in \mathbb{Z}$.

Lemma 2.7. [22] *Let M be a closed operator, $u \in L_{2\pi}^p(\mathbb{R}; [D(M)])$ and $u' \in L_{2\pi}^p(\mathbb{R}; X)$ for $1 \leq p < \infty$. Then, the following assertions are equivalent,*

(i) $\int_0^{2\pi} (Mu)'(t)dt = 0$ and there exist $x \in X$ such that $Mu(t) = x + \int_0^t (Mu)'(s)ds$ a.e. on $[0, 2\pi]$;

(ii) $\widehat{(Mu)'(k)} = ikM\hat{u}(k)$ for all $k \in \mathbb{Z}$.

Let a be a complex valued function. We define the set

$$\rho_{M,a}(A) = \{\lambda \in \mathbb{C} : (\lambda M - (\alpha + a(\lambda))A) : D(A) \cap D(B) \rightarrow X \\ \text{is invertible and } (\lambda M - (\alpha + a(\lambda))A)^{-1} \in \mathcal{B}(X)\},$$

and denote by $\sigma_{M,a}(A)$ the complementary set $\mathbb{C} \setminus \rho_{M,a}(A)$. If $M = I$, is the identity operator on X , $a \equiv 0$, and $\alpha = 1$, we denote simply the set $\rho_{M,a}(A)$ by $\rho(A)$ and as usual we call this set, the resolvent set of A . Denote by $\tilde{a}(\lambda)$ the Laplace transform of a . In what follows, we always assume that $\tilde{a}(ik)$ exists for all $k \in \mathbb{Z}$.

Henceforth, we use the following notation:

$$a_k := \tilde{a}(ik)$$

and we suppose that $a_k \neq -\alpha$ for all $k \in \mathbb{Z}$, where $\alpha \in \mathbb{R} \setminus \{0\}$.

Remark 2.8. Note that by the Riemann-Lebesgue lemma, the sequences $\{a_k\}_{k \in \mathbb{Z}}$ and $\{\frac{1}{\alpha + a_k}\}_{k \in \mathbb{Z}}$ ($\alpha \neq 0$) are bounded.

Finally, from [20] we recall the concept of n -regularity for $n = 1, 2, 3$. The general notion of n -regularity is the discrete analogue for the notion of n -regularity related to Volterra integral equations (see [23, Chapter I, Section 3.2]).

Definition 2.9. A sequence $\{c_k\}_{k \in \mathbb{Z}} \subset \mathbb{C} \setminus \{0\}$ is said to be:

(i) 1-regular, if the sequence $\left\{k \frac{(c_{k+1} - c_k)}{c_k}\right\}_{k \in \mathbb{Z}}$ is bounded.

(ii) 2-regular, if it is 1-regular and the sequence $\left\{k^2 \frac{(c_{k+1} - 2c_k + c_{k-1})}{c_k}\right\}_{k \in \mathbb{Z}}$ is bounded.

(iii) 3-regular, if it is 2-regular and the sequence $\left\{k^3 \frac{(c_{k+2} - 3c_{k+1} + 3c_k - c_{k-1})}{c_k}\right\}_{k \in \mathbb{Z}}$ is bounded.

Note that if $\{c_k\}_{k \in \mathbb{Z}}$ is 1-regular, then $\lim_{|k| \rightarrow \infty} c_{k+1}/c_k = 1$. For more details of n -regularity sequences, see [20].

3. MAXIMAL REGULARITY ON VECTOR-VALUED LEBESGUE SPACES

To characterize the maximal regularity, we begin with the study of the relation between multipliers and R -boundedness of a sequences of operators. Consider the problem

$$(3.1) \quad \begin{cases} \frac{d}{dt}(Mu(t)) = \alpha Au(t) + \int_{-\infty}^t a(t-s)Au(s)ds + f(t), & 0 \leq t \leq 2\pi, \\ Mu(0) = Mu(2\pi), \end{cases}$$

where $(A, D(A))$ and $(M, D(M))$ are closed linear operators on X , $D(A) \subseteq D(M)$, $a \in L^1(\mathbb{R}_+)$ is a scalar-valued kernel, $\alpha \in \mathbb{R} \setminus \{0\}$ and $f \in L^p_{2\pi}(\mathbb{R}, X)$, $p \geq 1$.

From [22], we recall that for a given closed operator M , and $1 \leq p < \infty$, the set $H^1_{per, M}(\mathbb{R}; [D(M)])$ is defined by

$$H^1_{per, M}(\mathbb{R}; [D(M)]) = \{u \in L^p_{2\pi}(\mathbb{R}; [D(M)]) : \exists v \in L^p_{2\pi}(\mathbb{R}; X), \hat{v}(k) = ikM\hat{u}(k) \text{ for all } k \in \mathbb{Z}\}.$$

When $M = I$, we denote $H^1_{per}(\mathbb{R}; X)$; see [3].

Lemma 3.1. *Let X be a UMD-space. Suppose that the sequence $\{a_k\}_{k \in \mathbb{Z}}$ is 1-regular. Then, for $\alpha \neq 0$, $\{\frac{1}{\alpha+a_k}I\}_{k \in \mathbb{Z}}$ is an L^p -multiplier.*

Proof. By Remarks 2.8 and 2.3 (g), $\{\frac{1}{\alpha+a_k}I\}_{k \in \mathbb{Z}}$ is R -bounded. Moreover,

$$\begin{aligned} k \left(\frac{1}{\alpha + a_{k+1}} - \frac{1}{\alpha + a_k} \right) &= k \left(\frac{a_k - a_{k+1}}{a_{k+1}} \right) \cdot \frac{a_{k+1}}{\alpha + a_{k+1}} \cdot \frac{1}{\alpha + a_k} \\ &= k \left(\frac{a_k - a_{k+1}}{a_{k+1}} \right) \cdot \frac{1}{\frac{\alpha}{a_{k+1}} + 1} \cdot \frac{1}{\alpha + a_k}. \end{aligned}$$

Since the 1-regularity of $\{a_k\}_{k \in \mathbb{Z}}$ imply the 1-regularity of $\{\frac{1}{a_k}\}_{k \in \mathbb{Z}}$, it follows that $\{k(\frac{a_k - a_{k+1}}{a_{k+1}})\}_{k \in \mathbb{Z}}$ is bounded and therefore $\{k(\frac{1}{\alpha+a_{k+1}} - \frac{1}{\alpha+a_k})I\}_{k \in \mathbb{Z}}$ is R -bounded. We conclude the proof of Lemma by Theorem 2.5. \blacksquare

The following Proposition is an extension of [22, Proposition 3.2].

Proposition 3.2. *Suppose that the sequence $\{a_k\}_{k \in \mathbb{Z}}$ is 1-regular. Let $A : D(A) \subseteq X \rightarrow X$ and $M : D(M) \subseteq X \rightarrow X$ be linear closed operators defined on a UMD space X . Suppose that $D(A) \subseteq D(M)$ and $\alpha \neq 0$. Then, the following assertions are equivalent:*

(i) $\{ik\}_{k \in \mathbb{Z}} \subset \rho_{M, \tilde{a}}(A)$ and $\{ikM(ikM - (\alpha + a_k)A)^{-1}\}_{k \in \mathbb{Z}}$ is an L^p -multiplier for $1 < p < \infty$;

(ii) $\{ik\}_{k \in \mathbb{Z}} \subset \rho_{M, \tilde{a}}(A)$ and $\{ikM(ikM - (\alpha + a_k)A)^{-1}\}_{k \in \mathbb{Z}}$ is R -bounded.

Proof. Define $N_k := (ikM - (\alpha + a_k)A)^{-1}$ and $M_k := ikM(ikM - (\alpha + a_k)A)^{-1}$, $k \in \mathbb{Z}$. By the Closed Graph Theorem we can show that if $ik \in \rho_{M, \tilde{a}}(A)$, then M_k are bounded operators for each $k \in \mathbb{Z}$. By Proposition 2.4 it follows that (i) implies (ii). Conversely, by Theorem 2.5 is sufficient to prove that the set $\{k(M_{k+1} - M_k)\}_{k \in \mathbb{Z}}$ is R -bounded. In fact, we note the following

$$\begin{aligned}
(3.2) \quad k[M_{k+1} - M_k] &= k[i(k+1)MN_{k+1} - ikMN_k] \\
&= ikMN_{k+1}[(k+1)[ikM - (\alpha + a_k)A] - \\
&\quad k[i(k+1)M - (\alpha + a_{k+1})A]]N_k \\
&= ikMN_{k+1}[k(a_{k+1} - a_k)A - (\alpha + a_k)A]N_k \\
&= ikMN_{k+1}\left[k\frac{(a_{k+1} - a_k)}{\alpha + a_k}\right](\alpha + a_k)AN_k - M_k(\alpha + a_k)AN_k.
\end{aligned}$$

Since $\{a_k\}_{k \in \mathbb{Z}}$ is 1-regular and $\alpha \neq 0$, the sequence $\{k\frac{(a_{k+1} - a_k)}{\alpha + a_k}\}_{k \in \mathbb{Z}}$ is bounded. The identity $(\alpha + a_k)AN_k = M_k - I$, imply that $\{(\alpha + a_k)AN_k\}_{k \in \mathbb{Z}}$ is R -bounded. We conclude the proof using the Remark 2.3. \blacksquare

Next, we introduce the following definition of solution of (3.1).

Definition 3.3. We say that a function $u \in H_{per, M}^{1,p}(\mathbb{R}; [D(M)])$ is a strong L^p -solution of (3.1) if $u(t) \in D(A)$ and equation (3.1) holds for a.e. $t \in [0, 2\pi]$.

The following is the main result of this section. For $a \in L^1(\mathbb{R}_+)$ and A a closed operator, denote by $F(t) := (a * Au)(t) = \int_{-\infty}^t a(t-s)Au(s)ds$. An easy computation show that $\hat{F}(k) = A\tilde{a}(ik)\hat{u}(k)$, $k \in \mathbb{Z}$, where the hat $\hat{\cdot}$ denotes Fourier transform (see [18] for a proof of this assertion).

Theorem 3.4. Let X be a UMD space. Let $A : D(A) \subseteq X \rightarrow X$, $M : D(M) \subseteq X \rightarrow X$ be linear closed operators. Suppose that $D(A) \subseteq D(M)$, the sequence $\{a_k\}_{k \in \mathbb{Z}}$ is 1-regular and $(ikM - (\alpha + a_k)A)$ are closed operators for all $k \in \mathbb{Z}$ where $\alpha \neq 0$. Then, the following assertions are equivalent:

- (i) For every $f \in L_{2\pi}^p(\mathbb{R}, X)$, there exist a unique strong L^p -solution of (3.1);
- (ii) $\{ik\}_{k \in \mathbb{Z}} \subset \rho_{M, \tilde{a}}(A)$ and $\{ikM(ikM - (\alpha + a_k)A)^{-1}\}_{k \in \mathbb{Z}}$ is an L^p -multiplier for $1 < p < \infty$;
- (iii) $\{ik\}_{k \in \mathbb{Z}} \subset \rho_{M, \tilde{a}}(A)$ and $\{ikM(ikM - (\alpha + a_k)A)^{-1}\}_{k \in \mathbb{Z}}$ is R -bounded.

Proof. (i) \Rightarrow (ii). We follow the same lines of [3, Theorem 2.3] and [18]. Let $k \in \mathbb{Z}$ and $y \in X$. Define $f(t) = e^{ikt}y$. By hypothesis, there exists $u \in H_{per, M}^{1,p}(\mathbb{R}, [D(M)])$ such that $u(t) \in D(A)$ and $(Mu)'(t) = \alpha Au(t) + (a * Au)(t) + f(t)$. Taking Fourier transform on both sides, we have $\hat{u}(k) \in D(A)$ and,

$$\begin{aligned}
ikM\hat{u}(k) &= (\alpha + a_k)A\hat{u}(k) + \hat{f}(k) \\
&= (\alpha + a_k)A\hat{u}(k) + y.
\end{aligned}$$

Thus, $(ikM - (\alpha + a_k)A)\hat{u}(k) = y$ for all $k \in \mathbb{Z}$ and we conclude that $(ikM - (\alpha + a_k)A)$ is surjective. Let $x \in D(A)$. If $(ikM - (\alpha + a_k)A)x = 0$, then $u(t) = e^{ikt}x$ defines a periodic solution of (3.1). Hence $u \equiv 0$ by the assumption of uniqueness, and thus $x = 0$. Therefore, $(ikM - (\alpha + a_k)A)$ is bijective. We conclude, from the Closed Graph Theorem, that $ik \in \rho_{M, \hat{a}}(A)$ for all $k \in \mathbb{Z}$. We will see that $\{ikM(ikM - (\alpha + a_k)A)^{-1}\}_{k \in \mathbb{Z}}$ is an L^p -multiplier.

Using the Closed Graph Theorem, we have that there exist a constant $C > 0$ independent of $f \in L^p_{2\pi}(\mathbb{R}; X)$ such that

$$\|(Mu)'\|_{L^p} + \|a \star Au\|_{L^p} + \|\alpha Au\|_{L^p} \leq C\|f\|_{L^p}.$$

Note that for $f(t) = e^{itk}y, y \in X$, the solution u of (3.1) is given by $u(t) = (ikM - (\alpha + a_k)A)^{-1}e^{ikt}y$. Hence,

$$\|ikM(ikM - (\alpha + a_k)A)^{-1}y\| \leq C\|y\|.$$

We obtain that for $k \in \mathbb{Z}$, $ikM(ikM - (\alpha + a_k)A)^{-1}$ is a bounded operator. Let $f \in L^p_{2\pi}(\mathbb{R}, X)$, by hypothesis, there exist $u \in H^{1,p}_{per, M}(\mathbb{R}, [D(M)])$ such that $u(t) \in D(A)$ and $(Mu)'(t) = \alpha Au(t) + (a \star Au)(t) + f(t)$. Taking Fourier transform on both sides, and using that $(ikM - (\alpha + a_k)A)$ is invertible, we have $\hat{u}(k) \in D(A)$ and $\hat{u}(k) = (ikM - (\alpha + a_k)A)^{-1}\hat{f}(k)$. Now, since $u \in H^{1,p}_{per, M}(\mathbb{R}, [D(M)])$ and by definition of $H^{1,p}_{per, M}(\mathbb{R}, [D(M)])$, there exist $v \in L^p_{2\pi}(\mathbb{R}, X)$ such that $\hat{v}(k) = ikM\hat{u}(k)$ for all $k \in \mathbb{Z}$. Therefore, we have $\hat{v}(k) = ikM\hat{u}(k) = ikM(ikM - (\alpha + a_k)A)^{-1}\hat{f}(k)$.

(ii) \Rightarrow (i). Define $M_k = ikM(ikM - (\alpha + a_k)A)^{-1}$ and $N_k = (ikM - (\alpha + a_k)A)^{-1}$. Suppose that $\{ik\}_{k \in \mathbb{Z}} \subset \rho_{M, \hat{a}}(A)$ and $\{M_k\}_{k \in \mathbb{Z}}$ is an L^p -multiplier. For $f \in L^p_{2\pi}(\mathbb{R}, X)$ there exist $u \in L^p_{2\pi}(\mathbb{R}, X)$ such that $\hat{u}(k) = ikM(ikM - (\alpha + a_k)A)^{-1}\hat{f}(k)$, for all $k \in \mathbb{Z}$. The identity $I = M_k - (\alpha + a_k)AN_k$ imply that

$$\begin{aligned} \hat{u}(k) &= ikM(ikM - (\alpha + a_k)A)^{-1}\hat{f}(k) \\ &= (I + (\alpha + a_k)AN_k)\hat{f}(k). \end{aligned}$$

So, we obtain $\widehat{(u - f)}(k) = (\alpha + a_k)AN_k\hat{f}(k)$. By Lemma 3.1, the sequence $\{\frac{1}{\alpha + a_k}I\}_{k \in \mathbb{Z}}$ is an L^p -multiplier. Thus, for $u - f \in L^p_{2\pi}(\mathbb{R}, X)$ there exists $v \in L^p_{2\pi}(\mathbb{R}, X)$ such that $\hat{v}(k) = \frac{1}{\alpha + a_k}\widehat{(u - f)}(k) = AN_k\hat{f}(k)$. Since that $0 \in \rho_{M, \hat{a}}(A)$ we obtain that $A^{-1} \in \mathcal{B}(X)$, and therefore $w := A^{-1}v \in L^p_{2\pi}(\mathbb{R}, X)$ and $\hat{w}(k) = N_k\hat{f}(k)$. Hence $ikM\hat{w}(k) - (\alpha + a_k)A\hat{w}(k) = \hat{f}(k)$. Observe that for all $k \in \mathbb{Z}$, we have

$$\hat{u}(k) = ikM(ikM - (\alpha + a_k)A)^{-1}\hat{f}(k) = ikM\hat{w}(k).$$

Thus $w \in H^{1,p}_{per, M}(\mathbb{R}, [D(M)])$. Moreover $Mw(0) = Mw(2\pi)$, since $w(0) = w(2\pi)$ and $w(t) \in D(A)$. Since A and M are closed operators and $\widehat{(Mw)}'(k) = ikM\hat{w}(k) = (\alpha + a_k)A\hat{w}(k) + \hat{f}(k)$, for all $k \in \mathbb{Z}$, one has $(Mw)'(t) = \alpha Aw(t) + (a \star Au)(t) + f(t)$ a.e. by Lemmas 2.6 and 2.7. Thus, we conclude that w is a strong L^p -solution of (3.1).

Finally, to see the uniqueness, let $u \in H^{1,p}_{per, M}(\mathbb{R}, [D(M)])$ such that $(Mu)'(t) = \alpha Au(t) + (a \star Au)(t)$. Taking, again, Fourier transform on both sides we have $\hat{u}(k) \in D(A)$, and

$(ikM - (\alpha + a_k)A)\hat{u}(k) = 0$, for all $k \in \mathbb{Z}$. Since $(ikM - (\alpha + a_k)A)$ is invertible for all $k \in \mathbb{Z}$, we obtain $\hat{u}(k) = 0$ for all $k \in \mathbb{Z}$, and thus $u \equiv 0$.

(ii) \Leftrightarrow (iii). Proposition 3.2. ■

In the Hilbert case, we obtain a simple condition to existence and uniqueness of solutions of (3.1). Since in Hilbert spaces the concepts of R -boundedness and boundedness are equivalent [13], the proof of the next Corollary follows from Theorem 3.4.

Corollary 3.5. *Let H be a Hilbert space, $A : D(A) \subset H \rightarrow H$, and $M : D(M) \subset H \rightarrow H$ closed linear operators satisfying $D(A) \subseteq D(M)$ and $(ikM - (\alpha + a_k)A)$ are closed operators for all $k \in \mathbb{Z}$, where $\alpha \neq 0$. Then, for $1 < p < \infty$, the following assertions are equivalent:*

(i) *For every $f \in L^p_{2\pi}(\mathbb{R}, H)$, there exists a unique strong L^p -solution of (3.1);*

(ii) *$\{ik\}_{k \in \mathbb{Z}} \subset \rho_{M, \tilde{a}}(A)$ and $\sup_{k \in \mathbb{Z}} \|(ikM - (\alpha + a_k)A)^{-1}\| < \infty$.*

Note that the solution $u(\cdot)$ given in Theorem 3.4 satisfies the following maximal regularity property.

Corollary 3.6. *In the context of Theorem 3.4, if condition (iii) is fulfilled, we have $(Mu)', Au, a \star Au \in L^p_{2\pi}(\mathbb{R}, X)$. Moreover, there exists a constant $C > 0$ independent of $f \in L^p_{2\pi}(\mathbb{R}; X)$ such that*

$$(3.3) \quad \|(Mu)'\|_{L^p} + \|\alpha Au\|_{L^p} + \|a \star Au\|_{L^p} \leq C \|f\|_{L^p}.$$

Remark 3.7.

The Fejer's Theorem (see [3, Proposition 1.1]) can be used to construct the solution u given in the Theorem 3.4. More precisely, if $M_k = ikM(ikM - (\alpha + a_k)A)^{-1}$ satisfies the condition (ii) or (iii) in the Theorem 3.4, then for $f \in L^p_{2\pi}(\mathbb{R}, X)$, the solution $u \in L^p_{2\pi}(\mathbb{R}, X)$ of (3.1) is given by

$$u(\cdot) = \lim_{n \rightarrow \infty} \frac{1}{n+1} \sum_{m=0}^n \sum_{k=-m}^m e_k \otimes M_k \hat{f}(k),$$

where $e_k(t) := e^{ikt}$, $t \in \mathbb{R}$ and the convergence holds in $L^p_{2\pi}(\mathbb{R}, X)$.

4. MAXIMAL REGULARITY ON HÖLDER AND BESOV SPACES

In this section, we formulate analogous theorems to the above section, in the context of Hölder and Besov Spaces.

Besov spaces form one class of function spaces that are of special interest. The relatively complicated definition is recompensed by useful applications to differential equations (see [2] for a concrete model).

We consider solutions in $B^s_{p,q}((0, 2\pi); X)$, the vector-valued periodic Besov spaces for $1 \leq p \leq \infty, s > 0$. Is remarkable that in this case, there are no geometrical conditions on

the Banach space X . For the definition and main properties of these spaces we refer to [4] or [19]. For the scalar case, see [12], [24]. Contrary to the L^p case, the multiplier theorems established for vector-valued Besov spaces are valid for arbitrary Banach spaces X ; see [1], [4] and [17]. Note also that, for $0 < s < 1$, $B_{\infty,\infty}^s$ is the usual Hölder space C^s . We summarize some useful properties of $B_{p,q}^s((0, 2\pi); X)$. See [4, Section 2] for a proof.

- (i) $B_{p,q}^s((0, 2\pi); X)$ is a Banach space;
- (ii) If $s > 0$, then $B_{p,q}^s((0, 2\pi); X) \hookrightarrow L^p((0, 2\pi); X)$, and the natural injection from $B_{p,q}^s((0, 2\pi); X)$ into $L^p((0, 2\pi); X)$ is a continuous linear operator;
- (iii) Let $s > 0$. Then $f \in B_{p,q}^{s+1}((0, 2\pi); X)$ if and only if f is differentiable a.e. and $f' \in B_{p,q}^s((0, 2\pi); X)$.

We begin with the definition of operator valued Fourier multipliers in the context of periodic Besov spaces.

Definition 4.8. *Let $1 \leq p, q \leq \infty, s > 0$. A sequence $\{M_k\}_{k \in \mathbb{Z}} \subset \mathcal{B}(X, Y)$ is a $B_{p,q}^s$ -multiplier if for each $f \in B_{p,q}^s((0, 2\pi); X)$ there exists a function $g \in B_{p,q}^s((0, 2\pi); Y)$ such that*

$$\hat{g}(k) = M_k \hat{f}(k), \quad k \in \mathbb{Z}.$$

Definition 4.9. [20] *We say that $\{M_k\}_{k \in \mathbb{Z}} \subset \mathcal{B}(X, Y)$ is M -bounded if*

$$(4.4) \quad \sup_{k \in \mathbb{Z}} \|M_k\| < \infty, \quad \sup_{k \in \mathbb{Z}} \|k(M_{k+1} - M_k)\| < \infty,$$

$$(4.5) \quad \sup_{k \in \mathbb{Z}} \|k^2(M_{k+1} - 2M_k + M_{k-1})\| < \infty.$$

We recall the following operator-valued Fourier multiplier theorem on Besov spaces.

Theorem 4.10. [4] *Let X, Y be Banach spaces and let $\{M_k\}_{k \in \mathbb{Z}} \subseteq \mathcal{B}(X, Y)$ be a M -bounded sequence. Then for $1 \leq p, q \leq \infty, s \in \mathbb{R}$, $\{M_k\}_{k \in \mathbb{Z}}$ is an $B_{p,q}^s$ -multiplier.*

The following Proposition is the analogous of the Proposition 3.2.

Proposition 4.11. *Let $A : D(A) \subseteq X \rightarrow X, M : D(M) \subseteq X \rightarrow X$ be linear closed operators on a Banach space X . Suppose that $D(A) \subseteq D(M)$ and the sequence $\{a_k\}_{k \in \mathbb{Z}}$ is 2-regular and $\alpha \neq 0$. Then, the following assertions are equivalent*

(i) $\{ik\}_{k \in \mathbb{Z}} \subset \rho_{M, \bar{a}}(A)$ and $\{ikM(ikM - (\alpha + a_k)A)^{-1}\}_{k \in \mathbb{Z}}$ is an $B_{p,q}^s$ -multiplier for $1 \leq p, q \leq \infty$;

(ii) $\{ik\}_{k \in \mathbb{Z}} \subset \rho_{M, \bar{a}}(A)$ and $\sup_{k \in \mathbb{Z}} \|ikM(ikM - (\alpha + a_k)A)^{-1}\| < \infty$.

Proof. (i) \Rightarrow (ii). Follows the same lines as the proof in [18, Proposition 3.4]. (ii) \Rightarrow (i) For $k \in \mathbb{Z}$, define $M_k = ikM(ikM - (\alpha + a_k)A)^{-1}$ and $N_k = (ikM - (\alpha + a_k)A)^{-1}$. From the identity (3.2) we obtain:

$$(4.6) \quad \sup_{k \in \mathbb{Z}} \|k(M_{k+1} - M_k)\| < \infty,$$

proving (4.4). To verify (4.5), note that:

$$\begin{aligned}
k^2[M_{k+1} - 2M_k + M_{k-1}] &= ik^2M \left[(k+1)N_{k+1} - 2kN_k + (k-1)N_{k-1} \right] \\
&= ik^2MN_{k+1} \left[(k+1)N_k^{-1} - 2kN_{k+1}^{-1} + (k-1)N_{k+1}^{-1}N_{k-1}N_k^{-1} \right] N_k \\
&= ik^2MN_{k+1} \left[(k+1)N_k^{-1} - 2k \left[(ikM - (\alpha + a_k)A) - (a_{k+1} - a_k)A + iM \right] + \right. \\
&\quad \left. (k-1) \left[(i(k-1)M - (\alpha + a_{k-1})A) + 2iM - (a_{k+1} - a_{k-1})A \right] N_{k-1}N_k^{-1} \right] N_k \\
&= ik^2MN_{k+1} \left[(k+1)N_k^{-1} - 2kN_k^{-1} + 2k(a_{k+1} - a_k)A - 2ikM + \right. \\
&\quad \left. \left[(k-1)N_{k-1}^{-1} + 2i(k-1)M - (k-1)(a_{k+1} - a_{k-1})A \right] N_{k-1}N_k^{-1} \right] N_k \\
&= ik^2MN_{k+1} \left[(k+1)I - 2kI + 2k(a_{k+1} - a_k)AN_k - 2ikMN_k + \right. \\
&\quad \left. \left[(k-1)I + 2i(k-1)MN_{k-1} - (k-1)(a_{k+1} - a_{k-1})AN_{k-1} \right] \right] \\
&= ik^2MN_{k+1} \left[2k(a_{k+1} - a_k)AN_k - 2(M_k - M_{k-1}) - (k-1)(a_{k+1} - a_{k-1})AN_{k-1} \right] \\
&= ikMN_{k+1} \left[2k^2(a_{k+1} - a_k)AN_k - 2k(M_k - M_{k-1}) - k(k-1)(a_{k+1} - a_{k-1})AN_{k-1} \right] \\
&= ikMN_{k+1} \left[2k^2 \frac{(a_{k+1} - a_k)}{\alpha + a_k} (M_k - I) - 2k(M_k - M_{k-1}) - k(k-1) \frac{(a_{k+1} - a_{k-1})}{\alpha + a_{k-1}} (M_{k-1} - I) \right].
\end{aligned}$$

Using the identities,

$$2k^2(a_{k+1} - a_k) = k^2(a_{k+1} - 2a_k + a_{k-1}) + k^2(a_{k+1} - a_{k-1}),$$

and

$$k(k-1)(a_{k+1} - a_{k-1}) = k^2(a_{k+1} - a_{k-1}) - k(a_{k+1} - a_{k-1}),$$

we obtain

$$\begin{aligned}
&2k^2 \frac{(a_{k+1} - a_k)}{\alpha + a_k} [M_k - I] - k(k-1) \frac{(a_{k+1} - a_{k-1})}{\alpha + a_{k-1}} [M_{k-1} - I] \\
&= k^2 \frac{(a_{k+1} - 2a_k + a_{k-1})}{\alpha + a_k} [M_k - I] + k^2 \frac{(a_{k+1} - a_{k-1})}{\alpha + a_k} \times \\
&\quad \times \left[\frac{\alpha(M_k - M_{k-1}) + (a_k - a_{k-1})I + a_{k-1}M_k - a_kM_{k-1}}{\alpha + a_{k-1}} \right] + k \frac{(a_{k+1} - a_k)}{\alpha + a_{k-1}} [M_{k-1} - I] \\
&= k^2 \frac{(a_{k+1} - 2a_k + a_{k-1})}{\alpha + a_k} [M_k - I] + k \frac{(a_{k+1} - a_{k-1})}{\alpha + a_k} \left[\frac{\alpha}{\alpha + a_{k-1}} k(M_k - M_{k-1}) \right. \\
&\quad \left. + k \frac{(a_k - a_{k-1})}{\alpha + a_{k-1}} I + \frac{k}{\alpha + a_{k-1}} [a_{k-1}M_k - a_kM_{k-1}] \right] + k \frac{(a_{k+1} - a_k)}{\alpha + a_{k-1}} [M_{k-1} - I].
\end{aligned}$$

Because the identity

$$\begin{aligned}
\frac{k}{\alpha + a_{k-1}} [a_{k-1}M_k - a_kM_{k-1}] &= \frac{a_{k-1}}{\alpha + a_{k-1}} k[M_k - M_{k-1}] + k \frac{(a_{k-1} - a_k)}{\alpha + a_{k-1}} M_{k-1}, \\
k \frac{(a_{k+1} - a_{k-1})}{\alpha + a_k} &= k \frac{(a_{k+1} - a_k)}{\alpha + a_k} + \frac{k}{k-1} (k-1) \frac{(a_k - a_{k-1})}{a_{k-1}} a_{k-1} \frac{1}{\alpha + a_k},
\end{aligned}$$

$$k \frac{(a_{k+1} - a_k)}{a_{k-1}} = k \frac{(a_{k+1} - a_k)}{a_k} \frac{a_k}{\alpha + a_{k-1}},$$

and

$$k[M_k - M_{k-1}] = \frac{k}{k-1}(k-1)[M_k - M_{k-1}],$$

are valid, and from the fact that $\{k(M_{k+1} - M_k)\}_{k \in \mathbb{Z}}$ is bounded and $\{a_k\}_{k \in \mathbb{Z}}$ is 2-regular, we conclude from the above identities and Remark 2.8 that,

$$(4.7) \quad \sup_k \|k^2(M_{k+1} - 2M_k + M_{k-1})\| < \infty.$$

Thus, $\{M_k\}_{k \in \mathbb{Z}}$, is M -bounded and therefore, by Theorem 4.10, $\{M_k\}_{k \in \mathbb{Z}}$ is an $B_{p,q}^s$ -multiplier. \blacksquare

Lemma 4.12. *Let X be a Banach space. Suppose that the sequence $\{a_k\}_{k \in \mathbb{Z}}$ is 2-regular. Then, for $\alpha \neq 0$, $\{\frac{1}{\alpha+a_k}I\}_{k \in \mathbb{Z}}$ is an $B_{p,q}^s$ -multiplier for $1 \leq p, q \leq \infty$.*

Proof. Define $m_k := \frac{1}{\alpha+a_k}$, $k \in \mathbb{Z}$. By Remark 2.8, the sequence $\{m_k\}_{k \in \mathbb{Z}}$ is bounded. Moreover, $\{m_k\}_{k \in \mathbb{Z}}$ satisfies the identities,

$$k[m_{k+1} - m_k] = -k \frac{a_{k+1} - a_k}{\alpha + a_k} \frac{1}{\alpha + a_{k+1}},$$

and

$$\begin{aligned} k^2[m_{k+1} - 2m_k + m_{k-1}] &= -\frac{\alpha}{(\alpha + a_{k+1})(\alpha + a_k)(\alpha + a_{k-1})} k^2(a_{k+1} - 2a_k + a_{k-1}) + \\ &\quad \frac{2}{(\alpha + a_{k+1})(\alpha + a_k)(\alpha + a_{k-1})} k(a_k - a_{k-1})k(a_{k+1} - a_k) - \\ &\quad \frac{a_k}{(\alpha + a_{k+1})(\alpha + a_k)(\alpha + a_{k-1})} k^2(a_{k+1} - 2a_k + a_{k-1}). \end{aligned}$$

Since $\{a_k\}_{k \in \mathbb{Z}}$ is 2-regular, we conclude that $\{m_k\}_{k \in \mathbb{Z}}$ is M -bounded and therefore $\{m_k\}_{k \in \mathbb{Z}}$ is an $B_{p,q}^s$ -multiplier. \blacksquare

Definition 4.13. *Let $1 \leq p, q \leq \infty$ and $s > 0$. A function $u \in B_{p,q}^s((0, 2\pi); [D(A)])$ is said to be a strong $B_{p,q}^s$ -solution of (3.1) if $Mu \in B_{p,q}^{s+1}((0, 2\pi); X)$ and equation (3.1) holds for a.e. $t \in (0, 2\pi)$.*

The next Theorem, is the main result of this section and is the analogous version of Theorem 3.4 in the context of Besov spaces. We remark that there are no special conditions in the space X .

Theorem 4.14. *Let $1 \leq p, q \leq \infty$ and $s > 0$. Let $A : D(A) \subseteq X \rightarrow X$, $M : D(M) \subseteq X \rightarrow X$ be linear closed operators on a Banach space X . Suppose that $D(A) \subseteq D(M)$, the sequence $\{a_k\}_{k \in \mathbb{Z}}$ is 2-regular and $(ikM - (\alpha + a_k)A)$ are closed operators for all $k \in \mathbb{Z}$ where $\alpha \neq 0$. Then, the following assertions are equivalent*

(i) For every $f \in B_{p,q}^s((0, 2\pi); X)$ there exist a unique strong $B_{p,q}^s$ -solution of (3.1);

(ii) $\{ik\}_{k \in \mathbb{Z}} \subset \rho_{M, \tilde{a}}(A)$ and $\{ikM(ikM - (\alpha + a_k)A)^{-1}\}_{k \in \mathbb{Z}}$ is an $B_{p,q}^s$ -multiplier for $1 \leq p, q \leq \infty$;

(iii) $\{ik\}_{k \in \mathbb{Z}} \subset \rho_{M, \tilde{a}}(A)$ and $\sup_{k \in \mathbb{Z}} \|ikM(ikM - (\alpha + a_k)A)^{-1}\| < \infty$.

Proof.

(i) \Rightarrow (iii). Suppose that for every $f \in B_{p,q}^s((0, 2\pi); X)$ there exist a unique strong $B_{p,q}^s$ -solution of (3.1). Fix $x \in X$ and $k \in \mathbb{Z}$. Define $f(t) = e^{itk}x$. Then $f \in B_{p,q}^s((0, 2\pi); X)$. By hypothesis there exist $u \in B_{p,q}^s((0, 2\pi); [D(A)])$ with $Mu \in B_{p,q}^{s+1}((0, 2\pi); X)$ such that $u(t) \in D(A)$ and $(Mu)'(t) = \alpha Au(t) + (a \star Au)(t) + f(t)$ a.e. $t \in (0, 2\pi)$. By Lemma 2.7 we have $ikM\hat{u}(k) = \alpha A\hat{u}(k) + a_k A\hat{u}(k) + x$. Following the same reasoning in the proof of Theorem 3.4 we obtain that $(ikM - (\alpha + a_k)A)$ is invertible for all $k \in \mathbb{Z}$. Let $M_k := ikM(ikM - (\alpha + a_k)A)^{-1}$. We will see that $\{M_k\}_{k \in \mathbb{Z}}$ is bounded. Using the Closed Graph Theorem, we have that there exist a constant C independent of f such that

$$\|Mu\|_{B_{p,q}^{s+1}((0, 2\pi); X)} + \|\alpha Au\|_{B_{p,q}^s((0, 2\pi); [D(A)])} + \|a \star Au\|_{B_{p,q}^s((0, 2\pi); [D(A)])} \leq C\|f\|_{B_{p,q}^s((0, 2\pi); X)}.$$

Note that for $f(t) = e^{itk}x$, the solution u of (3.1) is given by $u(t) = (ikM - (\alpha + a_k)A)^{-1}e^{ikt}x$. Hence,

$$\sup_{k \in \mathbb{Z}} \|ikM(ikM - (\alpha + a_k)A)^{-1}x\| \leq C\|x\|.$$

(ii) \Rightarrow (i). Define $M_k = ikM(ikM - (\alpha + a_k)A)^{-1}$ and $N_k = (ikM - (\alpha + a_k)A)^{-1}$. Suppose that $\{ik\}_{k \in \mathbb{Z}} \subset \rho_{M, \tilde{a}}(A)$ and $\{M_k\}_{k \in \mathbb{Z}}$ is an $B_{p,q}^s$ -multiplier. For $f \in B_{p,q}^s((0, 2\pi); X)$ there exist $u \in B_{p,q}^s((0, 2\pi); X)$ such that $\hat{u}(k) = ikM(ikM - (\alpha + a_k)A)^{-1}\hat{f}(k)$, for all $k \in \mathbb{Z}$. The identity $I = M_k - (\alpha + a_k)AN_k$ imply that

$$\begin{aligned} \hat{u}(k) &= ikM(ikM - (\alpha + a_k)A)^{-1}\hat{f}(k) \\ &= (I + (\alpha + a_k)AN_k)\hat{f}(k). \end{aligned}$$

So, we obtain $\widehat{(u - f)}(k) = (\alpha + a_k)AN_k\hat{f}(k)$. By Lemma 4.12, the sequence $\{\frac{1}{\alpha + a_k}I\}_{k \in \mathbb{Z}}$ is an $B_{p,q}^s$ -multiplier. Thus, for $u - f \in B_{p,q}^s((0, 2\pi); X)$ there exists $v \in B_{p,q}^s((0, 2\pi); X)$ such that $\hat{v}(k) = \frac{1}{\alpha + a_k}\widehat{(u - f)}(k) = AN_k\hat{f}(k)$. Since that $0 \in \rho_{M, \tilde{a}}(A)$ we obtain that $A^{-1} \in \mathcal{B}(X)$, and therefore $w := A^{-1}v \in B_{p,q}^s((0, 2\pi); X)$ and $\hat{w}(k) = N_k\hat{f}(k)$. Hence $ikM\hat{w}(k) - (\alpha + a_k)A\hat{w}(k) = \hat{f}(k)$. Observe that for all $k \in \mathbb{Z}$, we have

$$\hat{u}(k) = ikM(ikM - (\alpha + a_k)A)^{-1}\hat{f}(k) = ikM\hat{w}(k).$$

Thus, by uniqueness of Fourier coefficients, $u(t) = (Mw)'(t)$. Since $u \in B_{p,q}^s((0, 2\pi); X)$, then $(Mw)' \in B_{p,q}^s((0, 2\pi); X)$ and therefore, $Mw \in B_{p,q}^{s+1}((0, 2\pi); [D(A)])$. Moreover $Mw(0) = Mw(2\pi)$, since $w(0) = w(2\pi)$ and $w(t) \in D(A)$.

Since A and M are closed operators and $\widehat{(Mw)'(k)} = ikM\hat{w}(k) = (\alpha + a_k)A\hat{w}(k) + \hat{f}(k)$, for all $k \in \mathbb{Z}$, one has $(Mw)'(t) = \alpha Aw(t) + (a \star Au)(t) + f(t)$ a.e. by Lemmas 2.6 and 2.7.

We conclude that $w \in B_{p,q}^s((0, 2\pi); X)$ is a strong $B_{p,q}^s$ -solution to (3.1).

Finally, to see the uniqueness, let $u \in B_{p,q}^s((0, 2\pi); [D(A)])$ such that $(Mu)'(t) = \alpha Au(t) + (a \star Au)(t)$. Taking, again, Fourier transform on both sides we have $\hat{u}(k) \in D(A)$, and $(ikM - (\alpha + a_k)A)\hat{u}(k) = 0$, for all $k \in \mathbb{Z}$. Since $(ikM - (\alpha + a_k)A)$ is invertible for all $k \in \mathbb{Z}$, we obtain $\hat{u}(k) = 0$ for all $k \in \mathbb{Z}$, and thus $u \equiv 0$.

(iii) \Leftrightarrow (ii). Follows from Proposition 4.11. \blacksquare

5. MAXIMAL REGULARITY ON TRIEBEL-LIZORKIN SPACES

In this section, we study the existence and uniqueness of solutions to (3.1) in the context of Triebel-Lizorkin spaces; $F_{p,q}^s((0, 2\pi); X)$, where X is a Banach space, $1 \leq p, q \leq \infty$ and $s \in \mathbb{R}$. More details of these spaces can be found in [7] and the references therein.

The next definition and theorem are the analogous versions mentioned in the above sections.

Definition 5.15. *Let $1 \leq p, q \leq \infty, s \in \mathbb{R}$. A sequence $\{M_k\}_{k \in \mathbb{Z}} \subset \mathcal{B}(X, Y)$ is a $F_{p,q}^s$ -multiplier if for each $f \in F_{p,q}^s((0, 2\pi); X)$ there exists a function $g \in F_{p,q}^s((0, 2\pi); Y)$ such that*

$$\hat{g}(k) = M_k \hat{f}(k), \quad k \in \mathbb{Z}.$$

We recall the following result due to [7].

Theorem 5.16. [7] *Let X, Y be Banach spaces and let $\{M_k\}_{k \in \mathbb{Z}} \subseteq \mathcal{B}(X, Y)$. Assume that*

$$(5.8) \quad \sup_{k \in \mathbb{Z}} \|M_k\| < \infty, \quad \sup_{k \in \mathbb{Z}} \|k(M_{k+1} - M_k)\| < \infty,$$

$$(5.9) \quad \sup_{k \in \mathbb{Z}} \|k^2(M_{k+1} - 2M_k + M_{k-1})\| < \infty,$$

$$(5.10) \quad \sup_{k \in \mathbb{Z}} \|k^3(M_{k+2} - 3M_{k+1} + 3M_k - M_{k-1})\| < \infty,$$

where $1 \leq p < \infty$ and $1 \leq q \leq \infty, s \in \mathbb{R}$, $\{M_k\}_{k \in \mathbb{Z}}$. Then $\{M_k\}_{k \in \mathbb{Z}}$ is an $F_{p,q}^s$ -multiplier.

Remark 5.17.

We remark that, if X, Y are *UMD* spaces in the above theorem, then the conditions (5.8) and (5.9) are sufficient for $\{M_k\}_{k \in \mathbb{Z}}$ to be an $F_{p,q}^s$ -multiplier.

The definition of solution of the equation (3.1) in the Triebel-Lizorkin spaces is the same that in the Besov case. The proof of following theorem is similar to Theorem 4.14. We omit the details.

Theorem 5.18. *Let $1 \leq p, q \leq \infty$ and $s > 0$. Let $A : D(A) \subseteq X \rightarrow X$, $M : D(M) \subseteq X \rightarrow X$ be linear closed operators on a Banach space X . Suppose that $D(A) \subseteq D(M)$, the sequence $\{a_k\}_{k \in \mathbb{Z}}$ is 3-regular and $(ikM - (\alpha + a_k)A)$ are closed operators for all $k \in \mathbb{Z}$ where $\alpha \neq 0$. Then, the following assertions are equivalent*

- (i) For every $f \in F_{p,q}^s((0, 2\pi); X)$ there exist a unique strong $F_{p,q}^s$ -solution of (3.1);
- (ii) $\{ik\}_{k \in \mathbb{Z}} \subset \rho_{M, \tilde{a}}(A)$ and $\{ikM(ikM - (\alpha + a_k)A)^{-1}\}_{k \in \mathbb{Z}}$ is an $F_{p,q}^s$ -multiplier for $1 \leq p, q \leq \infty$;
- (iii) $\{ik\}_{k \in \mathbb{Z}} \subset \rho_{M, \tilde{a}}(A)$ and $\sup_{k \in \mathbb{Z}} \|ikM(ikM - (\alpha + a_k)A)^{-1}\| < \infty$.

6. APPLICATIONS

We conclude the paper, with some applications of our results.

Example 6.19.

Let us consider the boundary value problem

$$(6.11) \quad \frac{\partial(m(x)u(t, x))}{\partial t} - \Delta u = \int_{-\infty}^t a(t-s)\Delta u(s, x)ds + f(t, x), \quad \text{in } [0, 2\pi] \times \Omega$$

$$(6.12) \quad u = 0, \quad \text{in } [0, 2\pi] \times \partial\Omega,$$

where Ω is a bounded domain in \mathbb{R}^n with a smooth boundary $\partial\Omega$, $m(x) \geq 0$ is a given measurable bounded function on Ω and f is a function on $[0, 2\pi] \times \Omega$.

Let M be the multiplication operator by m . If we take $X = H^{-1}(\Omega)$ then by [5, p.38] (see also references therein), we have that there exists a constant $c > 0$ such that

$$\|M(zM - \Delta)^{-1}\| \leq \frac{c}{1 + |z|},$$

whenever $\operatorname{Re} z \geq -c(1 + |\operatorname{Im}(z)|)$. Thus, the inequality

$$\|ikM(ikM - ((1 + a_k)\Delta)^{-1}\| = \frac{|k|}{|1 + a_k|} \left\| M \left(\frac{ik}{1 + a_k} M - \Delta \right)^{-1} \right\| \leq c,$$

holds, if $\operatorname{Re} \left(\frac{ik}{1 + a_k} \right) \geq -c(1 + |\operatorname{Im} \left(\frac{ik}{1 + a_k} \right)|)$, for all $k \in \mathbb{Z}$, that is, if

$$(6.13) \quad k\beta_k \geq -c((1 + \alpha_k)^2 + \beta_k^2 + |k(1 + \alpha_k)|),$$

is valid for all $k \in \mathbb{Z}$, where α_k and β_k denotes the real and imaginary part of a_k , respectively. In particular, if $a(t) := \frac{t^{b-1}}{\Gamma(b)}$, with b an odd integer, then one can check that $\beta_k = 0$ for all $k \in \mathbb{Z}$, thus the inequality (6.13) holds. Therefore, by Theorem 4.14 (or Corollary 3.5), we conclude that that for all $f \in L_{2\pi}^p(\mathbb{R}, H^{-1}(\Omega))$ there exists a unique solution for (6.11)-(6.12).

Example 6.20.

Consider, for $t \in [0, 2\pi]$ and $x \in [0, \pi]$, the problem

$$(6.14) \quad \frac{\partial}{\partial t} \left(\frac{\partial^2}{\partial x^2} + 1 \right) u(t, x) = -b \frac{\partial^2}{\partial x^2} u(t, x) - cu(t, x) +$$

$$(6.15) \quad \int_{-\infty}^t a(t-s) \left(b \frac{\partial^2}{\partial x^2} + c \right) u(s, x) ds + f(t, x)$$

$$(6.16) \quad u(t, 0) = u(t, \pi) = \frac{\partial^2}{\partial x^2} u(t, 0) = \frac{\partial^2}{\partial x^2} u(t, \pi) = 0$$

$$(6.17) \quad \left(\frac{\partial^2}{\partial x^2} + 1 \right) u(0, x) = \left(\frac{\partial^2}{\partial x^2} + 1 \right) u(2\pi, x),$$

where b is positive constant and $-2b < c < 4b$. If we take $X = C_0([0, \pi]) = \{u \in C([0, \pi]) : u(0) = u(\pi)\}$ and K the realization of $\frac{\partial^2}{\partial x^2}$ with domain

$$D(K) = \{u \in C^2([0, \pi]) : u(0) = u(\pi) = \frac{\partial^2}{\partial x^2} u(0) = \frac{\partial^2}{\partial x^2} u(\pi) = 0\},$$

then we take $M = K + I$, $A = bM + (c - b)I$. By [5, p.39, Ex.1.2] we have that:

$$\|M(zM - A)^{-1}\| \leq \frac{d}{1 + |z|}$$

for all $\operatorname{Re} z \geq -d(1 + |\operatorname{Im}(z)|)$, and d being a suitable positive constant. Therefore, as in the above example, if for all $k \in \mathbb{Z}$, the inequality

$$(6.18) \quad k\beta_k \geq -d((\alpha_k - 1)^2 + \beta_k^2 + |k(\alpha_k - 1)|),$$

is valid, then for all $f \in B_{p,q}^s((0, 2\pi), C_0([0, \pi]))$, $s > 0$, $1 \leq p, q \leq \infty$, by Theorem 4.14, the problem (6.14)-(6.17) has a unique strong solution u with regularity $\frac{\partial^2 u}{\partial x^2} \in B_{p,q}^s((0, 2\pi), C_0([0, \pi]))$. In particular, if $a(t) := e^{\gamma t}$, where $\gamma \in \mathbb{R}$, we can check that the inequality (6.18) holds with $d = 1$.

REFERENCES

- [1] H. Amann, *Operator-valued Fourier multipliers, vector-valued Besov spaces and applications*, Math. Nachr. **186** (1997), 5-56.
- [2] H. Amann, *On the strong solvability of the Navier-Stokes equations*, J. Math. Fluid. Mech. **2** (2000), 16-98.
- [3] W. Arendt, S. Bu, *The operator-valued Marcinkiewicz multiplier theorem and maximal regularity*, Math. Z. **240** (2002), 311-343.
- [4] W. Arendt, S. Bu, *Operator-valued Fourier multiplier on periodic Besov spaces and applications*, Proc. Edim. Math. Soc. **47** (2) (2004), 15-33.
- [5] V. Barbu, A. Favini, *Periodic problems for degenerate differential equations*, Rend. Instit. Mat. Univ. Trieste XXVIII (Suppl.) (1997) 29-57.

- [6] J. Bourgain, *Some remarks on Banach spaces in which martingale differences sequences are unconditional*, Arkiv Math. **21** (1983), 163-168.
- [7] S. Bu, J.M. Kim, *Operator-valued Fourier multipliers on periodic Triebel spaces*, Acta Math. Sinica (Engl. Ser.) **21** (2005), 1049-1056.
- [8] S. Bu, Y. Fang, *Maximal regularity for integro-differential equation on periodic Triebel-Lizorkin spaces*. Taiwanese J. Math. **12** (2) (2008), 281-292.
- [9] D.L. Burkholder, *A geometrical condition that implies the existence of certain singular integrals on Banach-space-valued functions*. In: Proc. Conference on Harmonic Analysis in Honor of Antoni Zygmund, Chicago 1981, pp.270-286, Wadsworth, Belmont, CA, 1983.
- [10] D.L. Burkholder, *Martingales and singular integrals in Banach spaces*. In: Handbook of the Geometry of Banach Spaces, Vol 1, pp.233-269, North-Holland, Amsterdam, (2001).
- [11] T.A. Burton, B. Zhang, *Periodic solutions of abstract differential equations with infinite delay*. J. Differential Equations, **90** (1991), 357-396.
- [12] P.L. Butzer, H. Berens, *Semi-groups of operators and approximation*, Die Grundlehren der Mathematische Wissenschaften, **145**, Springer Verlag, 1967.
- [13] R. Denk, M. Hieber, J. Prüss, *R-boundedness, Fourier multipliers and problems of elliptic and parabolic type*, Mem. Amer. Math. Soc. **166** (788), 2003.
- [14] A. Favini, A. Yagi, *Degenerate differential equations in Banach spaces*, Pure and Applied Math., **215**, Dekker, New York, Basel, Hong-Kong, 1999.
- [15] A. Favaron, A. Favini, *Maximal time regularity for degenerate evolution integro-differential equations*. J. Evol. Equ. **10** (2) (2010), 377-412.
- [16] M. Girardi, L. Weis, *Criteria for R-boundedness of operator families*, Lecture Notes in Pure and Appl. Math., **234**, Dekker, New York, 2003, 203-221.
- [17] M. Girardi, L. Weis, *Operator-valued Fourier multiplier theorems on Besov spaces*, Math. Nachr., **251**, (2003), 34-51.
- [18] V. Keyantuo, C. Lizama, *Fourier multipliers and integro-differential equations in Banach spaces*, J. London Math. Soc. **69** (3) (2004), 737-750.
- [19] V. Keyantuo, C. Lizama, *Maximal regularity for a class of integro-differential equations with infinite delay in Banach spaces*, Studia Math. **168** (1) (2005), 25-50.
- [20] V. Keyantuo, C. Lizama and V. Pobleto, *Periodic solutions of integro-differential equations in vector-valued function spaces*, J. Differential Equations, **246** (3) (2009), 1007-1037.
- [21] J. Lagnese, *Boundary Stabilization of Thin Plates*, SIAM, Philadelphia, (1989).

- [22] C. Lizama, R. Ponce, *Periodic solutions of degenerate differential equations in vector-valued function spaces*. *Studia Math.*, **202** (2011), 49-63.
- [23] J. Prüss. *Evolutionary Integral Equations and Applications*. *Monographs Math.*, **87**, Birkhäuser Verlag, 1993.
- [24] H.J.-Schmeisser, H. Triebel, *Topics in Fourier Analysis and Function Spaces*, Chichester, Wiley, 1987.
- [25] L. Weis, *A new approach to maximal L_p -regularity*. *Lect. Notes Pure Appl. Math.* **215** Marcel Dekker, New York, (2001), 195-214.

UNIVERSIDAD DE SANTIAGO DE CHILE, DEPARTAMENTO DE MATEMÁTICA, FACULTAD DE CIENCIAS,
CASILLA 307-CORREO 2, SANTIAGO-CHILE.

E-mail address: carlos.lizama@usach.cl

E-mail address: rodrigo.ponce@usach.cl